Section Editor
Mitchell S.V. Elkind,
MD, MS

Teaching *NeuroImages*: Acute generalized suppression on continuous EEG heralds clinical and radiologic deterioration

Tadeu Fantaneanu, MD
Vincent Alvarez, MD
Jong Woo Lee, MD, PhD

Correspondence to
Dr. Lee:
Jlee38@partners.org

A 79-year-old woman was admitted to the hospital following 2 generalized tonic clonic seizures and a large right intracerebral hemorrhage (ICH). She deteriorated 72 hours later, losing her right pupillary reflex. A CT scan at that time showed stable ICH volume with effacement of the right suprasellar cistern, concerning for impending herniation. Interestingly, her continuous EEG (cEEG) showed a marked change 90 minutes prior to her clinical deterioration, characterized by acute generalized suppression of cerebral activity (figure). cEEG changes in patients with raised intracranial pressure vary by etiology, ranging from lateralized periodic discharges in the acute phase of injury to bifrontally predominant generalized rhythmic delta activity to suppression of EEG activity. cEEG changes may precede clinical and radiographic deterioration and should be followed closely in patients with elevated intracranial pressure (ICP).¹ Identification of cEEG changes in critically ill patients with changes in ICP should be treated with greater clinical vigilance and may lead to early interventions such as hyperosmolar therapy or even surgical interventions such as hematoma evacuation or decompressive craniectomy.²

AUTHOR CONTRIBUTIONS

Dr. Fantaneanu contributed to the design/conceptualization of this study, interpretation of the data, and the drafting/revision of the manuscript. Dr. Alvarez contributed to the design/conceptualization of this study, interpretation of the data, and the drafting/revision of the manuscript. Dr. Lee contributed to the design/conceptualization of this study, interpretation of the data, and the drafting/revision of the manuscript. Dr. Lee has access to all the data and takes responsibility for the data, accuracy of the data analysis, and the conduct of this research.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

T. Fantaneanu and V. Alvarez report no disclosures relevant to the manuscript. J. Lee has received research funding from UCB, Inc., the Duke Clinical Research Institute, and Sunovion, Inc. He is a consultant for SleepMed/DigiTrace. Go to Neurology.org for full disclosures.

REFERENCES

1. Newey CR, Sarwal A, Hantus S. Continuous electroencephalography (cEEG) changes precede clinical changes in a case of progressive cerebral edema. *Neurocrit Care* 2013;18:261–265.
2. Perez-Barcena J, Llopart-Pou JA, O'Phelan KH. Intracranial pressure monitoring and management of intracranial hypertension. *Crit Care Clin* 2014;30:735–750.

Download teaching slides:
[Neurology.org](#)

From the Department of Neurology (T.F., V.A., J.W.L.), Brigham and Women's Hospital, Boston, MA; and Hôpital du Valais (V.A.), Sion, Switzerland.

Figure Timeline of radiographic and cEEG monitoring record with spectral analysis

(A) Admission CT: right intracerebral hemorrhage; continuous EEG (cEEG): paradoxical left lateralized periodic discharges and intermixed rhythmic delta activity as well as bilateral, but right greater than left, hemispheric slowing and attenuation. (B) CT at time of clinical change; cEEG: suppression. (C) Diffuse edema and herniation 9 hours later. (D) EEG spectral analysis (8 hours) shows loss of cerebral activity 1.5 hours before clinical change (T0).

Neurology®

Teaching *NeuroImages*: Acute generalized suppression on continuous EEG heralds clinical and radiologic deterioration

Tadeu Fantaneanu, Vincent Alvarez and Jong Woo Lee

Neurology 2015;84:e119-e120

DOI 10.1212/WNL.0000000000001490

This information is current as of April 20, 2015

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/84/16/e119.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2015/04/19/WNL.0000000000001490.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/84/16/e119.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): EEG http://n.neurology.org/cgi/collection/eeg EEG; see Epilepsy/Seizures http://n.neurology.org/cgi/collection/eeg_see_epilepsy-seizures Video/ EEG use in epilepsy http://n.neurology.org/cgi/collection/video__eeg_use_in_epilepsy
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2015 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.