- pathogenesis of experimental murine cerebral malaria. *J Immunol* 1996;15:1620–1624.
23. Chimelli L, Mahler–Araujo MB. Fungal infections. *Brain Pathol* 1997;7:613–627.
 24. Graninger W, Prada J, Neifer S, Zotter G, Thalhammer F, Kramsner P. Upregulation of ICAM-1 by *Plasmodium falciparum*: in vitro and in vivo studies. *J Clin Pathol* 1994;47:653–655.
 25. Jakobsen P, Morris–Jones S, Ronn A, et al. Increased plasma concentrations of sICAM-1, sVCAM-1 and sELAM-1 in patients with *Plasmodium falciparum* or *P. vivax* malaria and association with disease severity. *Immunol* 1994;83:655–659.
 26. Jander S, Heidenreich F, Stoll G. Serum and CSF levels of soluble intercellular adhesion molecule-1 (ICAM-1) in inflammatory neurologic diseases. *Neurology* 1993;43:1809–1813.
 27. Wenisch C, Parschalk B, Burgmann H, Looareesuwan S,

- Graninger W. Decreased serum levels of TGF-beta in patients with acute *Plasmodium falciparum* malaria. *J Clin Immunol* 1995;15:69–73.
28. Flanders KC, Ren RF, Lipka CF. Transforming growth factor-betas in neurodegenerative disease. *Prog Neurobiol* 1998;54:71–85.
 29. Brown H, Turner G, Rogerson S, et al. Cytokine expression in the brain in human cerebral malaria. *J Infect Dis* 1999;180:1762–1746.
 30. Yong VW, Krekoski CA, Forsyth PA, Bell R, Edwards DR. Matrix metalloproteinases and diseases of the CNS. *Trends Neurosci* 1998;21:75–80.
 31. Paul R, Lorenzi S, Koedel U, et al. Matrix metalloproteinases contribute to the blood–brain barrier disruption during bacterial meningitis. *Ann Neurol* 1998;44:592–600.

NeuroImages

Figure. Diffusion-weighted MRI study. Arrows indicate areas of acute ischemic injury.

Recurrent silent cerebral embolism

A.E. Baird, FRACP, PhD, S. Warach, MD, PhD, Bethesda, MD

Multiple silent cerebral ischemic lesions in space and time occurred in a 58-year-old man with an untreated left ventricular cardiac thrombus.^{1–2} This patient initially presented with a sudden onset of expressive dysphasia, right-sided weakness and hypotension, with a history of myocardial infarction and poor left ventricular function. The first MR diffusion-weighted imaging (DWI) study was obtained two hours after the onset of symptoms. The hyperintense areas (arrows) indicate acute ischemic injury. DWI lesions were present in the left frontal region, the right cerebral hemisphere, and the left cerebellar hemisphere. On serial MR studies new clinically silent small

lesions were detected. At 26 hours and 8 days new lesions had developed in the right cerebral hemisphere (arrows). The left frontal lesion had shown some signs of lesion evolution with hyperintensity and some enlargement. At 75 days a new lesion was present adjacent to the posterior horn of the right lateral ventricle. An apical thrombus was subsequently found on transesophageal echocardiography. A previous bleeding episode had precluded anticoagulation until this time.

1. Baird AE, Lovblad KO, Schlaug G, Edelman RR, Warach S. Multiple acute stroke syndrome. Marker of embolic disease? *Neurology* 2000;54:674–678.
2. Babikian VL, Caplan LR. Brain embolism is a dynamic process with variable characteristics. *Neurology* 2000;54:797–801.

Neurology®

Recurrent silent cerebral embolism

Neurology 2000;55;111

DOI 10.1212/WNL.55.1.111

This information is current as of July 12, 2000

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/55/1/111.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.