Correction

Safety and efficacy of pallidal or subthalamic nucleus stimulation in advanced Parkinson’s disease

In the article “Safety and efficacy of pallidal or subthalamic nucleus stimulation in advanced Parkinson’s disease” by Volkmann et al. (*Neurology* 2001;56:548–551), table 2 was printed incorrectly. In table 2, the Unified PD Rating Scale (UPDRS) motor score (off medication/on stimulation) at 12 months after surgery should be 26 ± 15.46 , not 16.7 ± 11.64 , which is the UPDRS motor score on medication/on stimulation. The authors apologize for this error. The corrected table 2 is printed below.

Table 2 Motor scores, activities of daily living (ADL) scores, and levodopa-equivalent daily dose (LEDD) before and after surgery in internal pallidum (GPi)-stimulated patients

Rating scale	Score range	Off-period score, mean $\pm$ SD			On-period score, mean $\pm$ SD		
		Baseline	Stimulation on		Baseline	Stimulation on	
			6 mo	12 mo		6 mo	12 mo
UPDRS ADL	0–52	21.0 $\pm$ 6.72	9.5 $\pm$ 7.80*	12.1 $\pm$ 8.52†	12.10 $\pm$ 7.10	5.9 $\pm$ 7.05†	5.8 $\pm$ 6.18†
UPDRS motor	0–108	52.5 $\pm$ 14.16	22.9 $\pm$ 15.48*	26 $\pm$ 15.46*	30.2 $\pm$ 18.50	15.2 $\pm$ 11.74*	16.7 $\pm$ 11.64†
Schwab & England	0–100	51.8 $\pm$ 15.37	81.8 $\pm$ 9.82*	78.9 $\pm$ 12.69†	71.8 $\pm$ 17.22	89.1 $\pm$ 8.31†	90 $\pm$ 8.66†
Dyskinesia Rating Scale	0–4	—	—	—	2.0 $\pm$ 1.10	0.4 $\pm$ 0.51§	0.4 $\pm$ 0.53§
UPDRS subscores							
Bradykinesia	0–4	2.5 $\pm$ 0.70	1.2 $\pm$ 1.01*	1.3 $\pm$ 0.68†	1.5 $\pm$ 0.69	0.8 $\pm$ 0.73†	1.3 $\pm$ 0.68†
Rigidity	0–4	2.2 $\pm$ 0.65	0.8 $\pm$ 0.64†	1.1 $\pm$ 0.94†	1.4 $\pm$ 0.98	0.6 $\pm$ 0.51†	0.7 $\pm$ 0.73†
Tremor	0–4	0.8 $\pm$ 0.79	0.1 $\pm$ 0.19*	0.2 $\pm$ 0.24†	0.3 $\pm$ 0.65	0.1 $\pm$ 0.20†	0.1 $\pm$ 0.11†
Posture & gait	0–4	2.2 $\pm$ 0.72	1.2 $\pm$ 0.90*	0.8 $\pm$ 0.69†	1.1 $\pm$ 0.72	0.6 $\pm$ 0.60†	0.7 $\pm$ 0.81†
Speech & swallowing	0–4	1.3 $\pm$ 0.56	0.8 $\pm$ 0.79†	0.9 $\pm$ 0.74†	0.8 $\pm$ 0.70	0.6 $\pm$ 0.74†	0.4 $\pm$ 0.52†
Dyskinesias	0–4	—	—	—	1.2 $\pm$ 1.00	0.4 $\pm$ 0.43*	0.2 $\pm$ 0.40*
LEDD, mg		—	—	—	836 $\pm$ 391	700 $\pm$ 311†	605 $\pm$ 189†

p Values calculated by two-tailed Wilcoxon test.

* *p* < 0.005; †*p* = NS; §*p* < 0.001.

UPDRS = Unified PD Rating Scale.

Neurology®

Safety and efficacy of pallidal or subthalamic nucleus stimulation in advanced Parkinson's disease

Neurology 2001;57;1354
DOI 10.1212/WNL.57.7.1354

This information is current as of October 9, 2001

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/57/7/1354.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.