Mitoxantrone-induced heart failure in MS

Ghalie et al. reviewed records of 1,378 patients who received mitoxantrone in MS clinical trials. At median cumulative dose 62.5 mg/m² and follow-up 29 months, 2 of 1,378 patients (<0.20%) experienced congestive heart failure (CHF). Whether the incidence of CHF in MS patients will prove to be higher with greater cumulative mitoxantrone doses is not yet established.

see page 909

Fatal leukemia with mitoxantrone treatment of MS

Brassat et al. report acute leukemia in a mitoxantrone-treated patient with severe MS—the first case among 802 French mitoxantrone-treated MS patients.

see page 954

Mitoxantrone-related toxicity in multiple sclerosis

Commentary by John W. Rose, MD

Mitoxantrone is an important new therapy recently approved by the FDA for treatment of severe relapsing and secondary progressive multiple sclerosis. In this issue of *Neurology*, two articles discuss potential side effects of the drug. The article by Ghalie et al. examines the cardiotoxicity of the drug in multiple sclerosis patients demonstrating a low incidence of clinical congestive heart failure across several studies. The result of a 0.2% incidence for heart failure is much lower than the incidence in patients treated for neoplasias, a group frequently treated with other cardiotoxic

therapies prior to mitoxantrone administration. It is clear, however, that careful monitoring of cardiac function is important in the evaluation of patients treated even with low-dose mitoxantrone.¹

Of concern is the report by Brassat et al. of chemotherapy-related acute myeloblastic leukemia following mitoxantrone therapy. While leukemia occurred in only one patient of over 800 patients receiving a full course of mitoxantrone in the study, this is the second case of leukemia reported in a multiple sclerosis patient treated with the drug.² These manuscripts on mitox-

antrone-related toxicity underscore the requirement of appropriate patient selection and monitoring.

References

1. Strotman J, Spindler M, Weilbach F, Gold R, Ertl G, Wolfram V. Myocardial function in patients with multiple sclerosis treated with low-dose mitoxantrone. *Am J Cardiol* 2002;89:1222–1225.
2. Vicari A, Ciceri F, Folli F, Lanzi R, Colombo B, Comi G, Camba L. Acute promyelocytic leukemia following mitoxantrone as single agent for the treatment of multiple sclerosis. *Leukemia* 1998;12:441–442.

Diffusion MR imaging of neonates with brain injury

McKinstry et al. prospectively characterized changes in water diffusion that take place during the first week after brain injury at birth in 10 term infants. The sensitivity of diffusion MR imaging for detecting injury varied markedly with time after injury, being most sensitive between 2 and 4 days. During the first day after injury, diffusion MR imaging misses up to 30% of injuries in term infants. Thus, a “normal” diffusion MR scan obtained during the first day of life should be interpreted with caution.

see page 824

The accompanying editorial by Melhem notes the challenge that this study posed—obtaining serial MRI studies in critically ill newborns—and considers the study's limitations. Whether diffusion MR will be able to demonstrate changes in a larger spectrum of neonatal disorders, such as those with hypoxia from respiratory compromise, or in premature infants, can only be determined with further studies.

see page 798

continued on page 796

Arterial reocclusion after IV TPA

Alexandrov and Grotta used TCD to study patients with MCA occlusion treated with IV TPA. After initial partial or complete recanalization, reocclusion was observed in 34% of these patients. Reocclusion accounted for two-thirds of cases with neurologic deterioration following improvement and was associated with higher mortality.

see page 862

NSAIDs reduce AD incidence but . . .

Zandi et al. observed a reduced incidence of AD among users of NSAIDs in their 3-year follow-up assessment of the prospectively studied Cache County population. However, their data suggest a "lag" of several years between NSAID use and prevention of AD onset. This "lag" suggests that NSAIDs would not prevent progression from mild cognitive impairment in AD.

see page 880

Poverty and dementia: Cause or effect?

In a population-based study Anttila et al. show that dementia is associated with low-income level at old age and a decrease in income level from midlife. They note that the low old age income level is more likely to reflect the effects of the disease than the reverse.

see page 887

Migraine and impaired cognitive function

In a longitudinal birth cohort study done over a 23-year period Waldie et al. found that migraineurs have impaired verbal functioning and lessened academic success. However, the deficits are unrelated to duration of headache history, and are likely to be neurodevelopmental in origin rather than due to repeated vascular insults.

see page 904

New pathomechanism of muscular dystrophy

A study of a patient with Ullrich congenital muscular dystrophy by Ishikawa et al. suggests that the loss of a link between interstitium and basal lamina may be a new molecular pathomechanism of muscular dystrophy.

see page 920

continued on page 797

On-line video: Denny-Brown's treatment of Wilson's disease with BAL

In 1951, Denny-Brown and Porter reported that Wilson's disease could be treated effectively with BAL. This first demonstration that treatment of a biochemical disorder could alleviate dyskinesias, even in adults, is documented by Vilensky et al. in the films of Denny-Brown and the five initial patients—illustrating this historical landmark in neurology.

see page 914

Carotid stenting for dissecting aneurysms

Binaghi et al. report stent angioplasty in two patients with spontaneous dissecting aneurysms of the cervical internal carotid artery. Repeat angiography at 1 year disclosed complete repair of the arteries.

see page 935

*(Left) Initial. Arrow—true lumen; hollow arrow—false channel.
(Right) One-year ICA.*

Neurogenic flare reaction in neuropathy

Bickel et al. analyzed the axon-reflex mediated flare reaction with laser Doppler scanning and found flare size but not intensity reduced in patients with suspected small fiber neuropathy.

see page 917

Neurology®

September 24 Highlights
Neurology 2002;59;795-797
DOI 10.1212/WNL.59.6.795

This information is current as of September 24, 2002

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/59/6/795.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/59/6/795.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/59/6/795.full##otherarticles
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.