would therefore be insufficient to sustain memory functions? Baxendale et al. also state, “fMRI techniques cannot be used to assess the unilateral capacity of individual temporal lobe structures.” It is a mistake to exclude fMRI from providing such information in the future.⁷ Although the Wada test provides more direct measurement of memory function in one hemisphere, fMRI activation can potentially provide an indirect assessment of memory reserves.

Bassel W. Abou-Khalil, MD, *Nashville, TN*;
Bradley L. Schlaggar, MD, PhD, *St. Louis, MO*

Copyright © 2003 by AAN Enterprises, Inc.

References

1. Abou-Khalil B, Schlaggar BL. Is it time to replace the Wada test? Editorial. *Neurology* 2002;59:160–161.
2. Baxendale S. The Role of Functional MRI in the Presurgical Investigation of Temporal Lobe Epilepsy Patients: A Clinical Perspective and Review. *Journal of Clinical and Experimental Neuropsychology* 2002;24:664–676.
3. Jayakar P, Bernal B, Santiago Medina L, et al. False lateralisation of language cortex on functional MRI after a cluster of focal seizures. *Neurology* 2002;58:490–492.
4. Baxendale S. Amnesia in temporal lobectomy patients: historical perspective and review. *Seizure* 1988;7:15–24.
5. Davies KG, Bell BD, Bush AJ, Wyler AR. Prediction of verbal memory loss in individuals after anterior temporal lobectomy. *Epilepsia* 1998;39:820–828.
6. Gaillard WD, Balsame L, Xu B, et al. Language dominance in partial epilepsy patients identified with an fMRI reading task. *Neurology* 2002;59:256–265.
7. Golby AJ, Poldrack RA, Illes J, Chen D, Desmond JE, Gabrieli JDE. Memory lateralization in medial temporal lobe epilepsy assessed by functional MRI. *Epilepsia* 2002;43:855–863.

Correction

Assessment of low-flow CSF drainage as a treatment for AD: Results of a randomized pilot study

In the article, “Assessment of low-flow CSF drainage as a treatment for AD: Results of a randomized pilot study” by Silverberg et al. (*Neurology* 2002;59:1139–1145), the authors’ equity interests were not adequately disclosed in the Acknowledgment portion of this article. Dr. Silverberg is on the Board of Directors of Eunoe, Inc. and Dr. Dawn McGuire, Dr. Elliott Levinthal, Dr. Edith Sullivan, Dr. Daniel Bloch, Dr. Martha Mayo, and Mr. Thomas Saul have an equity interest in Eunoe, Inc. Eunoe, Inc. was the corporate sponsor of the study and manufacturer of the “CogniShunt.” Full disclosure of this information was made in the cover letter that accompanied the manuscript submission and on the author disclosure forms. It was not the intent of the authors to keep this information from the reader and the authors apologize for this oversight.

Neurology[®]

Assessment of low-flow CSF drainage as a treatment for AD: Results of a randomized pilot study

Neurology 2003;60;355
DOI 10.1212/WNL.60.2.355

This information is current as of January 28, 2003

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/60/2/355.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.