Figure. (A) T2-weighted MR image performed 6 hours after the onset of symptoms demonstrates a subtle hyperintensity area in the midbrain. (B) Diffusion-weighted MR image performed at the same time clearly demonstrates a hyperintensity area around the superior cerebellar peduncle just below and medial to the red nucleus. (C) T2-weighted MR image performed 4 days after the onset demonstrates a hyperintensity in the corresponding area.

MRI of Claude's syndrome

*Hiroyuki Asakawa, MD, Kiyoyuki Yanaka, MD, PhD,
Tadao Nose, MD, PhD, Ibaraki, Japan*

A 74-year-old man presented with sudden onset of right oculomotor palsy with sparing of the pupil and marked ataxia of the left upper limb. In 1912, the French neurologist Henri Claude first described the syndrome of ipsilateral oculomotor nerve palsy with

contralateral asynergia that bears his name.¹ Claude syndrome is caused by a lesion of the ventromedial midbrain, resulting in the combination of an ipsilateral oculomotor palsy and contralateral ataxia. There have been few studies describing the MR appearance of lesions associated with this syndrome.² MR imaging clearly localized the lesion of Claude syndrome in this case (figure).

Address correspondence and reprint requests to Dr. Kiyoyuki Yanaka, Department of Neurosurgery, Institute of Clinical medicine, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan; e-mail: kyanaka@md.tsukuba.ac.jp

1. Claude H. Syndrome pedonculaire de la region du noyau rouge. *Rev Neurol (Paris)* 1912;23:311-313.
2. Broadley SA, Taylor J, Waddy HM, Thompson PD. The clinical and MRI correlate of ischaemia in the ventromedial midbrain: Claude's syndrome. *J Neurol* 2001;248:1087-1089.

Neurology®

MRI of Claude's syndrome

Hiroyuki Asakawa, Kiyoyuki Yanaka and Tadao Nose

Neurology 2003;61;575

DOI 10.1212/WNL.61.4.575

This information is current as of August 25, 2003

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/61/4/575.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/61/4/575.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

fMRI

<http://n.neurology.org/cgi/collection/fmri>

Infarction

<http://n.neurology.org/cgi/collection/infarction>

MRI

<http://n.neurology.org/cgi/collection/mri>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.