Figure 1. Temporal skull contusions and orbital hematomas after syncope.

VIDEO Syncope during EEG recording

Beat Schaer, MD; Stefan Osswald, MD; Peter Fuhr, MD; and David Leppert, MD, Basel, Switzerland

A 75-year-old man had several syncope with severe skull contusions (figure 1). These events were not preceded by vegetative symptoms such as nausea, dizziness, or palpitations. Holter-EKG was normal and an EEG was performed for suspected epilepsy, during which complete atrioventricular block without ventricular escape-rhythm occurred (figure 2A). Six seconds later, the EEG showed progressive slowing of background activity and voltage reduction, followed by electrocerebral silence (figure 2B); clinically the patient lost cervical muscle tone and became unconscious. Thirty-three seconds later, sinus-rhythm reoccurred, followed by the return of cerebral activity in reverse manner (figure 2C). The patient regained consciousness instantly and, when addressed by the technician, he was fully responsive (see video clip). A pacemaker was implanted and the patient was free of symptoms.

Clinical presentation of syncope does not often allow one to distinguish between cerebrovascular and epileptic origin. Despite normal 24-hour EKG, simultaneous EKG recording is mandatory when EEG is used in the evaluation of syncope.^{1,2}

1. Pitney MR, Beran RG, Jones A. A simultaneous electrocardiogram is important when electroencephalography is used in the evaluation of loss of consciousness. *Electroencephalogr Clin Neurophysiol* 1994;90:246–248.
2. Brenner RP. Electroencephalography in syncope. *J Clin Neurophysiol* 1997;14:197–209.

Address correspondence and reprint requests to Dr. Beat Schaer, Department of Cardiology, University Hospital, CH-4031 Basel, Switzerland; e-mail: bschaer@uhbs.ch

Figure 2. (A) Complete AV-block without ventricular escape rhythm during EEG. (B) EEG showing diffuse slowing and finally electrocerebral silence. (C) Reappearance of ventricular escape rhythm (arrows) and of background activity 7 seconds later. QRS complexes indicated by arrows.

Neurology[®]

Syncope during EEG recording

Beat Schaer, Stefan Osswald, Peter Fuhr, et al.

Neurology 2004;63;E11

DOI 10.1212/01.WNL.0000137025.35912.1C

This information is current as of September 13, 2004

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/63/5/E11.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2004/09/13/63.5.E11.DC2 http://n.neurology.org/content/suppl/2004/09/20/63.5.E11.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/63/5/E11.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Epilepsy/Seizures http://n.neurology.org/cgi/collection/all_epilepsy_seizures EEG; see Epilepsy/Seizures http://n.neurology.org/cgi/collection/eeg_see_epilepsy-seizures Syncope http://n.neurology.org/cgi/collection/syncope
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.