CORRECTION

Cholesterol level and symptomatic hemorrhagic transformation after ischemic stroke thrombolysis

In the article "Cholesterol level and symptomatic hemorrhagic transformation after ischemic stroke thrombolysis" by O.Y. Bang et al. (*Neurology* 2007;68:737–742), the figure is incorrect. The correct figure and legend appear below. The publisher apologizes for the error.

Figure Severity and type of hemorrhagic transformation after thrombolysis for ischemic stroke by low-dose lipoprotein cholesterol level and premorbid statin use.

Neurology[®]

CORRECTION

Neurology 2007;68;1547
DOI 10.1212/01.wnl.0000265366.71843.9e

This information is current as of April 30, 2007

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/68/18/1547.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.