cases with low AD Braak stages, similar to the findings by others,^{1,5} was higher (70%) than for patients with high Braak stages (22%).

These studies suggest a higher influence of concomitant neuritic AD pathology on phenotype and clinical diagnostic accuracy of DLB than cortical LB distribution. The pathogenic relationship between both pathologies and their impact on clinical features and natural history of DLB is unclear.

Kurt A. Jellinger, Vienna, Austria

Disclosure: The author reports no conflicts of interest.

Editor's Note: The authors of the article were offered the opportunity to respond but declined.

Copyright © 2008 by AAN Enterprises, Inc.

1. Weisman D, Cho M, Taylor C, Adame A, Thal LJ, Hansen LA. In dementia with Lewy bodies, Braak stage determines phenotype, not Lewy body distribution. *Neurology* 2007;69:356–359.
2. Jellinger KA, Wenning GK, Seppi K. Predictors of survival in dementia with Lewy bodies and Parkinson dementia. *Neurodegen Dis* 2008 (in press).
3. Braak H, Del Tredici K, Rub U, de Vos RA, Jansen Steur EN, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 2003; 24:197–211.
4. McKeith IG, Dickson DW, Lowe J, et al. Diagnosis and management of dementia with Lewy bodies: third report of the DLB Consortium. *Neurology* 2005;65:1863–1872.
5. Merdes AR, Hansen LA, Jeste DV, et al. Influence of Alzheimer pathology on clinical diagnostic accuracy in dementia with Lewy bodies. *Neurology* 2003;60:1586–1590.

CORRECTION

Is it time to use observational data to estimate treatment effectiveness in multiple sclerosis?

In the editorial “Is it time to use observational data to estimate treatment effectiveness in multiple sclerosis?” (*Neurology*® 2007;69:1478–1479) by M. Trojano, the sentence “The popular belief that only RCTs [randomized, controlled trials] produce trustworthy results and that all observational studies are misleading does a disservice to patient care, clinical investigation, and education of health care professionals” was not attributed appropriately to the original author (Concato et al., *N Engl J Med* 2000;342:1887–1892). The author apologizes for this omission due to an oversight.

Neurology[®]

CORRECTION

Neurology 2008;70;408

DOI 10.1212/01.wnl.0000308386.00055.ac

This information is current as of January 28, 2008

**Updated Information &
Services**

including high resolution figures, can be found at:
<http://n.neurology.org/content/70/5/408.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.