Teaching *NeuroImage*: MRI in multiple system atrophy “Hot cross bun” sign and hyperintense rim bordering the putamina

João Massano, MD
Fernando Costa, MD
Goreti Nadais, MD

Address correspondence and
reprint requests to Dr. João
Massano, Department of
Neurology, Hospital de São João,
Alameda Prof. Hernani Monteiro,
4200-319 Porto, Portugal
jmassano@med.up.pt

(A) Axial T2 image showing the classic “hot cross bun” sign in the pons; axial (B) and coronal (C) T2 images depicting a hyperintense rim bordering the lateral margin of both putamina (white arrows); (D) sagittal T2 sequence disclosing pontine and cerebellar atrophy, the latter also evident in A. Right side of images is left side of patient in A, B, and C.

A 59-year-old man developed progressive gait ataxia, dysarthria, sexual dysfunction, and Raynaud phenomenon over 2 years. On examination he also manifested bradykinesia and rigidity, postural and action tremor of the upper limbs, and orthostatic hypotension. Family history was negative. Brain MRI disclosed the “hot cross bun” sign typical of multiple system atrophy (MSA) (figure).^{1,2} This sign is probably related to degeneration of transverse pontocerebellar fibers, whereas the hyperintense rim bordering the putamina has been associated with neuronal loss, reactive microgliosis, and astroglia-

sis.² Glial cytoplasmic inclusions, neuronal inclusions, and demyelination have been linked to MSA, but the mechanisms involved require further clarification.² MRI has diagnostic potential in parkinsonism-plus syndromes; several typical findings like those depicted here have been described.^{1,2}

REFERENCES

1. Schrag A, Good CD, Miszkil K, et al. Differentiation of atypical parkinsonian syndromes with routine MRI. *Neurology* 2000;54:697–702.
2. Wenning GK, Colosimo C, Geser F, Poewe W. Multiple system atrophy. *Lancet Neurol* 2004;3:93–103.

From the Departments of Neurology (J.M., G.N.) and Neuroradiology (F.C.), Hospital de São João; and Faculty of Medicine (J.M.), University of Porto, Portugal.

Disclosure: The authors report no disclosures.

Neurology®

Teaching *NeuroImage*: MRI in multiple system atrophy: "Hot cross bun" sign and hyperintense rim bordering the putamina

João Massano, Fernando Costa and Goreti Nadais

Neurology 2008;71:e38

DOI 10.1212/01.wnl.0000327520.99034.28

This information is current as of October 6, 2008

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/71/15/e38.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/71/15/e38.full#ref-list-1>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://n.neurology.org/content/71/15/e38.full##otherarticles>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Gait disorders/ataxia

http://n.neurology.org/cgi/collection/gait_disorders_ataxia

MRI

<http://n.neurology.org/cgi/collection/mri>

Multiple system atrophy

http://n.neurology.org/cgi/collection/multiple_system_atrophy

Parkinson's disease/Parkinsonism

http://n.neurology.org/cgi/collection/parkinsons_disease_parkinsonism

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.