resolution is an important attribute. As Dr. Zhu noted, advances in US technology permit visualization of nerve structures that was not possible a decade ago. Finally, as our editorial and Dr. Zhu highlight, real-time dynamic scanning of nerves is a critical advantage over MRI. All these considerations further confirm the utility of US in the evaluation of peripheral nerve disease.

© 2013 American Academy of Neurology

1. Padua L, Hobson-Webb LD. Ultrasound as the first choice for peripheral nerve imaging? *Neurology* 2013;80:1626–1627.
2. Wells PN, Liang HD, Young TP. Ultrasonic imaging technologies in perspective. *J Med Eng Technol* 2011;35:289–299.
3. Liu F, Zhu J, Wei M, Bao Y, Hu B. Preliminary evaluation of the sural nerve using 22-MHz ultrasound: a new approach for evaluation of diabetic cutaneous neuropathy. *PLoS One* 2012;7:e32730.
4. Zhu J, Liu F, Li D, Shao J, Hu B. Preliminary study of the types of traumatic peripheral nerve injuries by ultrasound. *Eur Radiol* 2011;21:1097–1101.

CORRECTION

Clinical Reasoning: A 62-year-old man with right wrist drop

In the Resident & Fellow article “Clinical Reasoning: A 62-year-old man with right wrist drop” by G. Cirillo et al. (*Neurology*® 2013;81:e81–e84), there is an error in the corresponding author’s title, which should have read Prof. Tedeschi, as well as errors involving the figure. The published figure should have been split into 2 figures and the first y-axis label in figure 1, panel A, should have read “1st ankle – EDB.” See corrected figures with titles and legends below. The publisher regrets the errors.

Figure 1 Nerve conduction study findings of right deep peroneal nerve

Distal nerve stimulation at ankle from extensor digitorum brevis (EDB) resulted in a normal compound muscle action potential (CMAP); subsequent single proximal stimulus at fibular head showed a significant drop of CMAP amplitude and area, which was still evident at the second single distal stimulus. Amp. p-p = amplitude measured at peak to peak; Area p- = area of negative peak; Fib. head = fibular head.

Figure 2 Repetitive nerve stimulation test findings

3-Hz repetitive nerve stimulation of the right median to abductor pollicis brevis (R-APB) (A) and right facial to nasalis (R-Na) (B) muscles showed significant decrement in both muscles.

Author disclosures are available upon request (journal@neurology.org).

Neurology®

Clinical Reasoning: A 62-year-old man with right wrist drop

Neurology 2013;81;1645

DOI 10.1212/01.wnl.0000435794.37174.34

This information is current as of October 28, 2013

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/81/18/1645.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.