1. Tung CE. Education Research: changing practice: residents' adoption of the atraumatic lumbar puncture needle. *Neurology* 2013;80:e180–e182.
2. Armon C, Evans R. Addendum to assessment: prevention of post-lumbar puncture headaches. *Neurology* 2005;65:510–512.
3. Tung CE, So YT, Lanberg MG. Cost comparison between the atraumatic and cutting lumbar puncture needle. *Neurology* 2012;78:109–133.
4. Fuchs VR, Milstein A. The \$640 billion question: why does cost-effective care diffuse so slowly? *N Engl J Med* 2011;364:1985–1987.

EDITORIAL EXPRESSION OF CONCERN: Structural and functional brain connectivity in presymptomatic familial frontotemporal dementia

With regard to the research article “Structural and functional brain connectivity in presymptomatic familial frontotemporal dementia” by Dopper et al. (*Neurology*® 2013;80:814–823; published ahead of print February 6, 2013), we are publishing this expression of concern to alert readers that the authors have discovered an error in the analyses that may influence the results and are planning to resolve the issue with additional studies. We are waiting for the authors to complete the reanalysis and will update readers when the results are received.

Neurology®

EDITORIAL EXPRESSION OF CONCERN: Structural and functional brain connectivity in presymptomatic familial frontotemporal dementia

Neurology 2014;82:735 Published Online before print January 24, 2014

DOI 10.1212/WNL.0000000000000230

This information is current as of January 24, 2014

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/82/8/735.full>

Errata

An erratum has been published regarding this article. Please see [next page](#) or:
</content/82/17/1570.full.pdf>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

CORRECTION

Structural and functional brain connectivity in presymptomatic familial frontotemporal dementia

In the article “Structural and functional brain connectivity in presymptomatic familial frontotemporal dementia” by E.G.P. Dopper et al. (*Neurology*® 2013;80:814–823), there is an error in the DNA results in 6 subjects. An Editorial Expression of Concern was published (*Neurology* 2014;82:735) and the authors have reanalyzed the data. Reanalysis of data concluded that there was significantly reduced resting-state functional connectivity within the salience network and reduced structural connectivity in the uncinate fasciculus in presymptomatic carriers, compared with controls, in the absence of gray matter atrophy. These findings support the conclusion of the original article; see authors’ full correction below. The authors regret the error.

In our article published in the February 26, 2013, issue of *Neurology*, we reported alterations in structural and functional connectivity in carriers of *progranulin* (*GRN*) and *microtubule-associated protein tau* (*MAPT*) mutations before the onset of symptoms of frontotemporal dementia (FTD).¹ In the follow-up study of our cohort of 75 at-risk individuals, we discovered an erroneous interchange of DNA results in 4 subjects and incorrect laboratory results in 2 additional subjects in our baseline article. Therefore, we reanalyzed our baseline data to investigate to what extent this error affected our results.

Similar to the reported results, reanalysis shows no differences in neuropsychological results between carriers and controls, except for significantly worse performances on the Letter Digit Substitution Test in carriers compared with controls. Reported age correlations with Stroop III and Happé cartoon performances in carriers, but not controls, are still present. In addition, there is a significant correlation between Rivermead Behavioral Memory Test performances with higher age in carriers, whereas the correlation for Ekman faces is no longer significant.

As reported, carriers and controls show no statistical differences in gray matter volume, except for a small cluster of higher volume in the right precentral gyrus in carriers that emerged in the reanalysis.

The reported differences in fractional anisotropy and radial diffusivity in frontotemporal white matter tracts are no longer significant, but using the uncinate fasciculus and forceps minor, tracts most consistently affected in FTD,^{2–6} as regions of interest,⁷ we find significantly reduced fractional anisotropy in the right uncinate fasciculus in carriers compared with controls.

The main reported resting-state fMRI findings were (1) a correlation between reduced frontoinsula (FI) connectivity with the anterior midcingulate cortex (aMCC) and age in carriers but not controls, and (2) reduced aMCC connectivity with parietal regions in carriers compared with controls. Both findings are still present in our reanalysis. In addition, we now find significantly decreased aMCC connectivity with FI and surrounding structures in carriers compared with controls, which strengthens our conclusion that salience network functional connectivity deteriorates prior to symptom onset. We also found reduced FI connectivity with several temporal and occipital regions.

To conclude, reanalysis of our data after error correction showed significantly reduced resting-state functional connectivity within the salience network and reduced structural connectivity in the uncinate fasciculus in presymptomatic carriers compared with controls, in the absence of gray matter atrophy. These findings support the conclusion of our original article that resting-state fMRI and diffusion tensor imaging changes in FTD develop before symptom onset.

1. Dopper EG, Rombouts SA, Jiskoot LC, et al. Structural and functional brain connectivity in presymptomatic familial frontotemporal dementia. *Neurology* 2013;80:814–823.
2. Agosta F, Scola E, Canu E, et al. White matter damage in frontotemporal lobar degeneration spectrum. *Cereb Cortex* 2012;22:2705–2714.
3. Hornberger M, Geng J, Hodges JR. Convergent grey and white matter evidence of orbitofrontal cortex changes related to disinhibition in behavioural variant frontotemporal dementia. *Brain* 2011;134:2502–2512.
4. Matsuo K, Mizuno T, Yamada K, et al. Cerebral white matter damage in frontotemporal dementia assessed by diffusion tensor tractography. *Neuroradiology* 2008;50:605–611.
5. Whitwell JL, Avula R, Senjem ML, et al. Gray and white matter water diffusion in the syndromic variants of frontotemporal dementia. *Neurology* 2010;74:1279–1287.
6. Zhang Y, Schuff N, Du AT, et al. White matter damage in frontotemporal dementia and Alzheimer’s disease measured by diffusion MRI. *Brain* 2009;132:2579–2592.
7. Mori S, Wakana S, Nagae-Poetscher L, van Zijl P. *MRI Atlas of Human White Matter*. Amsterdam: Elsevier; 2005.