Mystery Case: Lafora periodic acid–Schiff inclusion bodies

Igor de Assis Franco, MD
Eliza Teixeira da Rocha,
MD
Hugo Almeida Chaves de
Resende, MD
Nathália Moura da Silva
Guércio, MD
Moisés Pereira Pinto, MD
Leopoldo Antônio Pires,
MD
Thiago Cardoso Vale,
MD

Correspondence to
Dr. Vale:
thiago.vale@ufjf.edu.br

A 16-year-old boy presented with a 5-year history of progressive cognitive decline and behavioral change followed by generalized tonic-clonic and myoclonic seizures refractory to many anticonvulsants (valproic acid, phenobarbital, clonazepam, and topiramate) and cerebellar ataxia a year later. The son of consanguineous parents, he had a family history of 2 cousins who both had epilepsy and died at the ages of 15 and 25 years. Laboratory screening tests, including lactic acid, fundus examination, brain neuroimaging, and CSF, were normal. Skin biopsy (figure) revealed periodic acid–Schiff-positive intracellular polyglucosan inclusion bodies in myoepithelial and sweat gland duct cells. These findings are typical of Lafora disease, a fatal autosomal recessive disorder caused by mutations in one of 2 known genes, both located at chromosome 6: *EPM2A*, which encodes the protein laforin, and *EPM2B*, which encodes the protein malin. Differential diagnosis must be made with other causes of progressive myoclonic epilepsies, most commonly Unverricht-Lundborg disease (Baltic myoclonus), myoclonus epilepsy and ragged-red fibers syndrome, neuronal ceroid lipofuscinosis, and type 1 sialidosis. There is no treatment, and therapy is mainly supportive and symptomatic.^{1,2}

AUTHOR CONTRIBUTIONS

All authors contributed equally to the study concept and design, acquisition of data, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript for important intellectual content, and administrative, technical, and material support.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

REFERENCES

1. Monaghan TS, Delanty N. Lafora disease: epidemiology, pathophysiology and management. *CNS Drugs* 2010;24:549–561.
2. Newton GA, Sanchez RL, Swedo J, Smith EB. Lafora's disease: the role of skin biopsy. *Arch Dermatol* 1987;123:1667–1669.

MYSTERY CASE RESPONSES

The Mystery Case series was initiated by the *Neurology*® Resident & Fellow Section to develop the clinical reasoning skills of trainees. Residency programs, medical student preceptors, and individuals were invited to use this Mystery Case as an educational tool. Responses were solicited through a group e-mail sent to the American Academy of

Figure Lafora bodies on skin biopsy

(A) Histopathologic examination of the skin biopsy taken from the axillary region reveals round-to-oval intracytoplasmic, periodic acid–Schiff-positive, diastase-resistant inclusions within the myoepithelial and acinar cells of sweat glands, corresponding to Lafora bodies (arrows). Periodic acid–Schiff stain, 200×. (B) Lafora bodies in a magnified view (arrows). Periodic acid–Schiff stain, 400×.

From the Neurology Service (I.d.A.F., E.T.d.R., H.A.C.d.R., M.P.P., L.A.P., T.C.V.) and Pathology Service (N.M.d.S.G.), University Hospital, Department of Internal Medicine, Faculty of Medicine, Federal University of Juiz de Fora (UFJF), Juiz de Fora, Minas Gerais, Brazil.

Neurology Consortium of Neurology Residents and Fellows and through social media.

All of the respondents correctly identified the Lafora bodies in the figure and concluded that the underlying diagnosis is Lafora disease. Eighty percentage of respondents accurately stated that this is an autosomal recessive disease associated with mutations in the *EPM2A* and *EPM2B* genes.

This case illustrates the value of a detailed family history in guiding the appropriate diagnostic tests for rare genetic disorders such as the progressive myoclonic epilepsies.

Andrew Schepmyer, MD

University of British Columbia, Vancouver, Canada

Neurology®

Mystery Case: Lafora periodic acid–Schiff inclusion bodies

Igor de Assis Franco, Eliza Teixeira da Rocha, Hugo Almeida Chaves de Resende, et al.

Neurology 2015;85:e130–e131

DOI 10.1212/WNL.0000000000002054

This information is current as of October 26, 2015

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/85/17/e130.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/85/17/e130.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Epilepsy/Seizures http://n.neurology.org/cgi/collection/all_epilepsy_seizures All Genetics http://n.neurology.org/cgi/collection/all_genetics All Medical/Systemic disease http://n.neurology.org/cgi/collection/all_medical_systemic_disease All Pediatric http://n.neurology.org/cgi/collection/all_pediatric
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2015 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.