

# Orolingual and abdominal angioedema post thrombolysis and thrombectomy


Charline Benoit, MD, Marie Cantier, MD, Christine Rodriguez-Régent, MD, Olivier Gout, MD, and Michael Obadia, MD

## Correspondence

Dr. Benoit  
charlinebenoit5@gmail.com

*Neurology*® 2018;90:140-141. doi:10.1212/WNL.0000000000004834

## Figure 1 Imaging



(A) Magnetic resonance angiography shows right proximal occlusion of the middle cerebral artery. (B) Diffusion-weighted axial MRI shows right deep middle cerebral artery infarct.

A 53-year-old man treated with an angiotensin-converting enzyme (ACE) inhibitor presented with left hemiplegia (NIH Stroke Scale 23). Magnetic resonance angiography revealed a right proximal middle cerebral artery occlusion (figure 1). Thirty minutes after thrombolysis, orolingual swelling was observed. Two hours later, just after mechanical thrombectomy was completed, he developed painful abdominal edema, which improved after icatibant administration (figure 2). Bradykinin-mediated angioedema can be a complication of recombinant tissue plasminogen activator, typically orolingual, with increased risk for patients on ACE inhibitors.<sup>1</sup> We speculate that the procedure of thrombectomy, acting as a mechanical stress,<sup>2</sup> could have been a triggering factor for the unusual presentation of abdominal edema.

## Author contributions

Dr. Benoit: study concept and design. Dr. Cantier: analysis and interpretation. Dr. Rodriguez-Régent: analysis and interpretation. Dr. Olivier Gout: analysis and interpretation. Dr. Obadia: study supervision.

## Study funding

No targeted funding reported.

## Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org](http://Neurology.org) for full disclosures.

## References

1. Myslimi F, Caparros F, Dequatre-Ponchelle N, et al. Orolingual angioedema during or after thrombolysis for cerebral ischemia. *Stroke* 2016;47:1825–1830.
2. Aygören-Pürsün E, Martínez Saguer I, Kreuz W, Klingebiel T, Schwabe D. Risk of angioedema following invasive or surgical procedures in HAE type I and II the natural history. *Allergy* 2013;68:1034–1039.

From Neurologie (C.B., M.C., O.G., M.O.), Fondation Rothschild; and Neuroradiologie Interventionnelle (C.R.-R.), Centre Hospitalier Sainte-Anne, Paris, France.

**Figure 2** Left asymmetric upper and lower lips angioedema and abdominal painful edema



Left asymmetric upper and lower lips angioedema, contralateral to right hemispheric ischemia (A), and abdominal painful edema (C), both regressing after icatibant administration (B, D).

## Visit the *Neurology*<sup>®</sup> Website at [Neurology.org/N](http://Neurology.org/N)

- More article-based content on home pages
- Streamlined menus and navigation
- Enhanced blog sections for specialty areas
- Same experience on desktop, tablet, and mobile devices
- Audio summaries of current issues
- Improved article reading experience; links more evident (pdf, analytics, social media)
- *Neurology*<sup>®</sup> *Clinical Practice* initiative “Practice Current” global surveys will be accessible across sites

 Find *Neurology*<sup>®</sup> on Facebook: <http://tinyurl.com/neurologyfan>

 Follow *Neurology*<sup>®</sup> on Twitter: <https://twitter.com/GreenJournal>

# Neurology<sup>®</sup>

## Orolingual and abdominal angioedema post thrombolysis and thrombectomy

Charline Benoit, Marie Cantier, Christine Rodriguez-Régent, et al.

*Neurology* 2018;90;140-141

DOI 10.1212/WNL.0000000000004834

**This information is current as of January 15, 2018**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://n.neurology.org/content/90/3/140.full">http://n.neurology.org/content/90/3/140.full</a>
<b>References</b>	This article cites 2 articles, 1 of which you can access for free at: <a href="http://n.neurology.org/content/90/3/140.full#ref-list-1">http://n.neurology.org/content/90/3/140.full#ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>All Cerebrovascular disease/Stroke</b> <a href="http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke">http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke</a> <b>Critical care</b> <a href="http://n.neurology.org/cgi/collection/critical_care">http://n.neurology.org/cgi/collection/critical_care</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.neurology.org/about/about_the_journal#permissions">http://www.neurology.org/about/about_the_journal#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://n.neurology.org/subscribers/advertise">http://n.neurology.org/subscribers/advertise</a>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

