30. Lhermitte J. Syndrome de la calotte pedunculaire: les troubles psycho-sensoriels dans les lésions du mesencéphale. *Rev Neurol* 1922;38:1359–1365.
31. Pessah MA, Roffwarg HP. Spontaneous middle ear muscle activity in man: a rapid eye movement sleep phenomenon. *Science* 1972;178:773–776.
32. Braun AR, Balkin TJ, Wesensten NJ, et al. Dissociated pattern of activity in visual cortices and their projections during human rapid eye movement sleep. *Science* 1998;279:91–95.
33. Jouvet M, Delorme F. Locus coeruleus et sommeil paradoxal. *C R Soc Biol* 1965;159:895–899.
34. Aldrich MS. Narcolepsy. *N Engl J Med* 1990;323:389–394.
35. Sakai K. Physiological properties and afferent connections of the locus coeruleus and adjacent tegmental neurons involved in the generation of paradoxical sleep in the cat. *Prog Brain Res* 1991;88:31–45.
36. Mahowald MW, Schenck CH. Dissociated states of wakefulness and sleep. *Neurology* 1992;42:44–52.
37. Freud S. *Die Traumdeutung*. Vienna: Frantz Deuticke, 1900.
38. Gillin JC, Wyatt RJ. Schizophrenia: perchance a dream? *Int Rev Neurobiol* 1975;17:297–342.
39. Howland RH. Sleep-onset rapid eye movement periods in neuropsychiatric disorders: implications for the pathophysiology of psychosis. *J Nerv Ment Dis* 1997;185:730–738.
40. Douglass AB, Hays P, Pazderka F, Russell JM. Florid refractory schizophrenia that turn out to be treatable variants of HLA-associated narcolepsy. *J Nerv Ment Dis* 1991;179:12–17.
41. Johnson LC, Burdick JA, Smith J. Sleep during alcohol intake and withdrawal in the chronic alcoholic. *Arch Gen Psychiatry* 1970;22:406–418.
42. Hobson JA. How the brain goes out of its mind. *Endeavour* 1996;20:86–89.

NeuroImages

Figure. (A) MRI of cervical spine shows large flow voids, representing a vertebral arterial–epidural venous fistula, extending to the C5-6 level with cord compression. (B) Left vertebral angiogram demonstrates a dilated, tortuous vertebral artery supplying an arteriovenous fistula, which drains into large epidural veins in the spinal canal at the C3-4 level.

Arteriovenous fistula in neurofibromatosis

Xiaoping Ma, MD, PhD, Marina Amitina, MD,
Todd D. Rozen, MD, Philadelphia, PA

A 32-year-old woman with a history of neurofibromatosis type I presented with a subacute onset of left hemiparesis, neck pain, and urinary retention. Neurologic examination demonstrated a left hemiparesis and bilateral diffuse hyperreflexia. A loud bruit was auscultated on the

back of her neck. MRI of the cervical spine and vertebral angiography were completed (figure, A and B).

Neurofibromatosis type I is associated with central and peripheral nervous system tumors, but also cerebrocervical arterial lesions such as arteriovenous malformations, cervical artery aneurysms, and arteriovenous fistulas.¹ Vertebral arteriovenous fistulas can present as a radiculomyelopathy with neck pain and associated cervical bruits.¹

1. Schievink WI, Piepgras DG. Cervical vertebral artery aneurysms and arteriovenous fistulae in neurofibromatosis type I: case reports. *Neurosurgery* 1991;29:760–765.

Neurology®

Arteriovenous fistula in neurofibromatosis

Neurology 2000;55;288

DOI 10.1212/WNL.55.2.288

This information is current as of July 25, 2000

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/55/2/288.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.