


Teaching NeurolImage: Subclavian steal syndrome

Sashank Prasad, MD
Scott E. Kasner, MD

VIDEO

Address correspondence and
reprint requests to Dr. Sashank
Prasad, Department of
Neurology, Hospital of the
University of Pennsylvania, 3600
Spruce St., 3 Gates Bldg.,
Philadelphia, PA 19104
sashank.prasad@uphs.upenn.edu

Figure Cerebral angiography



Cerebral angiography reveals anterograde flow in the right vertebral artery (A) and delayed, retrograde flow through the left vertebral (B). There is complete occlusion of the proximal left subclavian artery (C, video). There is no evidence of additional stenoses. Attempts to recanalize the subclavian artery with angioplasty via femoral and brachial artery approaches were unsuccessful.

A 65-year-old woman presented with recurrent episodic vertigo lasting hours, accompanied only by headache. While golfing, she experienced transiently blurred vision. She denied arm claudication. Blood pressure was 160/78 in the right arm and 140/65 in the left arm. The left radial pulse was diminished. Cranial nerves, strength, sensation, and coordination were normal. Maneuvers to provoke symptoms were not attempted. MR angiography revealed an occlusion in the proximal left subclavian artery. Digital subtraction angiography was performed (see figure and video).

Proximal occlusion of the subclavian artery

causes retrograde flow through the ipsilateral vertebral artery.¹ Insufficient compensation by other arteries may result in posterior circulation ischemia, as in this case. However, many cases of “subclavian steal” may be asymptomatic. In these cases, invasive therapy should be withheld.²

REFERENCES

1. Reivich M, Holling EH, Roberts B, Toole JF. Reversal of blood flow through the vertebral artery and its effect on cerebral circulation. *N Engl J Med* 1961;265:878–881.
2. Bornstein NM, Norris JW. Subclavian steal: a harmless haemodynamic phenomenon? *Lancet* 1986;2:303–305.

Neurology[®]

Teaching *NeuroImage*: Subclavian steal syndrome

Sashank Prasad and Scott E. Kasner

Neurology 2007;69;E1

DOI 10.1212/01.wnl.0000265599.57822.85

This information is current as of July 9, 2007

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/69/2/E1.full>

Supplementary Material

Supplementary material can be found at:
<http://n.neurology.org/content/suppl/2007/06/28/69.2.E1.DC1>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/69/2/E1.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

