

Teaching *NeuroImage*: Acute spastic monoplegia secondary to spinal epidural venous engorgement in pregnancy

Michael J. Franco
Martina Vendrame,
MD, PhD
Zulfi Haneef, MD,
MRCP
S. Ausim Azizi, MD,
PhD

Address correspondence and
reprint requests to Dr. Martina
Vendrame, Temple University
Hospital, Neurology Department,
3401 North Broad Street,
Parkinson Pavilion Suite 558,
Philadelphia, PA 19140
vendrame@temple.edu

A 20-year-old, 33-week-pregnant woman presented with 2 days of backache and left leg weakness. She denied sphincter or sensory disturbance. Examination revealed proximal left leg weakness, bilateral hyperreflexia, and extensor plantars. A cesarean section was performed 5 days later. Two days postpartum, neurologic examination results returned to normal. Spine/abdomen MRIs before (figure, A–C) and after (figure, D–F) are presented.


Increased blood flow to the epidural venous plexus occurs due to compression on the inferior vena cava (IVC) by the pregnant uterus or by increased abdominal pressure.¹ This may present as

acute spastic paraparesis, but may also be asymptomatic,² which can be explained by the severity of IVC compression. MRI is helpful in excluding cord pathology, assuring favorable outcome.

REFERENCES

1. Hirabayashi Y, Shimizu R, Fukuda H, Saitoh K, Igarashi T. Effects of the pregnant uterus on the extradural venous plexus in the supine and lateral positions, as determined by magnetic resonance imaging. *Br J Anaesth* 1997;78:317–319.
2. Takiguchi T, Yamaguchi S, Tezuka M, Furukawa N, Kitajima T. Compression of the subarachnoid space by the engorged epidural venous plexus in pregnant women. *Anesthesiology* 2006;105:848–851.

Figure Spine/abdomen MRIs pre- and postpartum



(A) T2-weighted MRI of thoracic and lumbar spine revealing prominent epidural and paravertebral vessels in the lower thoracic and lumbar spinal canal consistent with vascular congestion. (B) Transverse T2-weighted image of the cord at level of T12-L1 showing increased signal, likely representing cord edema (white arrow). (C) MRI of the abdomen highlights compression of the inferior vena cava (IVC) (white arrow). (D) Repeated T2-weighted MRI of thoracic and lumbar spine showing resolution of vascular congestion. (E) Repeated transverse T2-weighted image of the cord at level of T12-L1 showing absence of the previously noted hyperintensity (white arrow). (F) MRI of the abdomen showing resolution of the IVC compression (white arrow).

From Temple University, Philadelphia, PA.

Disclosure: The authors report no disclosures.

Neurology[®]

Teaching *NeuroImage*: Acute spastic monoplegia secondary to spinal epidural venous engorgement in pregnancy

Michael J. Franco, Martina Vendrame, Zulfi Haneef, et al.

Neurology 2008;71:e1

DOI 10.1212/01.wnl.0000316310.85132.b3

This information is current as of June 30, 2008

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/71/1/e1.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/71/1/e1.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord Clinical neurology examination http://n.neurology.org/cgi/collection/clinical_neurology_examination MRI http://n.neurology.org/cgi/collection/mri Prognosis http://n.neurology.org/cgi/collection/prognosis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

