

# Teaching NeuroImage: Microangiopathic complications in pseudoxanthoma elasticum

D. Renard, MD  
G. Castelnovo, MD  
L. Jeanjean, MD  
H. Perrochia, MD  
H. Brunel, MD  
P. Labauge, MD, PhD

Address correspondence and reprint requests to Dr. Dimitri Renard, Department of Neurology, CHU Nîmes, Hôpital Caremeau, Place du Pr Debré, 30029 Nîmes Cedex 4, France  
dimitrirenard@hotmail.com

A 43-year-old woman without vascular risk factors presented with acute right sensorimotor deficit. Her medical history revealed pseudoxanthoma elasticum (PXE), diagnosed at age 29 by characteristic findings on skin biopsy (performed because of progressive skin lesions) and funduscopy (figure 1), complicated by peripheral artery disease and vision impairment.


Brain MRI showed pontine lacunar infarcts and microbleeds (figure 2), concordant with earlier reported microangiopathic complications in PXE.<sup>1,2</sup>

PXE is an inherited disorder characterized by progressive calcification and degeneration of elastic fibers, primarily affecting skin, eyes, and gastrointestinal and cardiovascular system. Management includes treatment of complications and strict control of cardiovascular risk factors.

## REFERENCES


1. van den Berg JS, Hennekam RC, Cruysberg JR, et al. Prevalence of symptomatic intracranial aneurysm and ischaemic stroke in pseudoxanthoma elasticum. *Cerebrovasc Dis* 2000;10:315–319.
2. Pavlovic AM, Zidverc-Trajkovic J, Milovic MM, et al. Cerebral small vessel disease in pseudoxanthoma elasticum: three cases. *Can J Neurol Sci* 2005;32:115–118.

**Figure 1** Patient, skin histopathology, and eye fundus



(A) Redundant axillary skin folds demonstrate loss of elasticity. (B) Histopathology of the skin on orcein stain showing calcium deposits (arrow) and swollen, fragmented elastic fibers (arrowheads). (C) Fundus photography of the eye shows characteristic angioid streaks (arrowheads), and a large retinal scar (white arrow) due to a macular hemorrhage (black arrows).

**Figure 2** Brain MRI



(A) MRI showing pontine ischemic T2 hyperintensities and (B) multiple foci of low intensity in the pons on echogradients T2-weighted images consistent with microbleeds.

From the Departments of Neurology (D.R., G.C., P.L.), Ophthalmology (L.J.), and Anatomopathology (H.P.), CHU Nîmes, Hôpital Caremeau, Nîmes; and Department of Neuroradiology (H.B.), CHU Montpellier, Hôpital Gui de Chauiac, Montpellier, France.

*Disclosure:* The authors report no disclosures.

# Neurology<sup>®</sup>

## Teaching *NeuroImage*: Microangiopathic complications in pseudoxanthoma elasticum

D. Renard, G. Castelnovo, L. Jeanjean, et al.

*Neurology* 2008;71:e69

DOI 10.1212/01.wnl.0000335934.11120.83

**This information is current as of November 24, 2008**

### Updated Information & Services

including high resolution figures, can be found at:  
<http://n.neurology.org/content/71/22/e69.full>

### References

This article cites 2 articles, 0 of which you can access for free at:  
<http://n.neurology.org/content/71/22/e69.full#ref-list-1>

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
[http://www.neurology.org/about/about\\_the\\_journal#permissions](http://www.neurology.org/about/about_the_journal#permissions)

### Reprints

Information about ordering reprints can be found online:  
<http://n.neurology.org/subscribers/advertise>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

