


Teaching NeuroImages: MRI in fibrodysplasia ossificans progressiva

Shiva Kumar R., MD
Keerthiraj B., MD
Chandrasekhran
Kesavadas, MD

Address correspondence and
reprint requests to Dr. Shiva
Kumar R., Department of
Neurology, Sree Chitra Tirunal
Institute for Medical Sciences and
Technology, Trivandrum
695011, Kerala, India
drshivakumar1995@yahoo.co.in

Figure Radiologic findings



Skeletal X-rays show clinodactyly, short thumb, and metacarpals (A), and hallux valgus with short phalanx (B). The CT scan shows calcification around paraspinal muscles (C). T2-weighted MRI of the brain shows dentate nucleus hyperintensity (D); STIR-coronal MRI (E), T2-weighted (F), and postcontrast (G) show increased signal and contrast enhancement in the muscles and fascial planes of the thoracic wall and paraspinal muscles.

A 3-year-old boy presented with abnormal neck posturing and rapidly growing lumps over the neck. The presence of typical malformed great toes and ectopic calcification (figure, A–C) and an affected family member confirmed the diagnosis of fibrodysplasia ossificans progressiva (FOP).¹ MRI of the cervicothoracic region (figure, C, E–G) revealed diffuse hyperintense signals extending along the neck, thoracic wall, and paraspinal muscles. The presence of dentate nucleus hyperintensity on brain MRI indicates that FOP may not be purely a musculoskeletal disorder, but a multisystem disorder including CNS involvement (figure, D). There was pseudodystonia of the neck without any other neurologic signs. Because ec-

topic ossification might not be present in the early stages, diagnosis of FOP based on radiographs alone may be delayed. In the presence of characteristic skeletal malformations, MRI can detect prososeous lesions and avoid additional diagnostic procedures, including biopsy.²

REFERENCES

1. Kaplan FS, Xu M, Glaser DL, et al. Early diagnosis of fibrodysplasia ossificans progressiva. *Pediatrics* 2008;121:e1295–e1300.
2. Merchant R, Sainani NI, Lawande MA, Pungavkar SA, Patkar DP, Walawalkar A. Pre- and post-therapy MR imaging in fibrodysplasia ossificans progressiva. *Pediatr Radiol* 2006;36:1108–1111.

From the Departments of Neurology (S.K.R.) and Imaging Sciences and Interventional Radiology (K.B., C.K.), Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, Kerala, India.

Disclosure: The authors report no disclosures.

Neurology[®]

Teaching *NeuroImages*: MRI in fibrodysplasia ossificans progressiva

R. Shiva Kumar, B. Keerthiraj and Chandrasekhran Kesavadas

Neurology 2010;74:e20

DOI 10.1212/WNL.0b013e3181cef7d1

This information is current as of February 8, 2010

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/74/6/e20.full
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/74/6/e20.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://n.neurology.org/cgi/collection/all_genetics All Pediatric http://n.neurology.org/cgi/collection/all_pediatric Clinical neurology examination http://n.neurology.org/cgi/collection/clinical_neurology_examination MRI http://n.neurology.org/cgi/collection/mri Muscle disease http://n.neurology.org/cgi/collection/muscle_disease
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

