

Child Neurology: A patient with dissimilar eye color and deafness

Chetan R. Soni, MD, MS,
MHA
Gyanendra Kumar, MD

Address correspondence and
reprint requests to Dr. Chetan R.
Soni, Mason Eye Institute, One
Hospital Drive, Columbia,
MO 65212
sonic@health.missouri.edu

Waardenburg syndrome (WS) is a rare genetic disorder with developmental anomalies of tissues derived from the neural crest and characterized by auditory and pigmentary findings. Failure of neural crest-derived melanocyte differentiation results in a spectrum of phenotypic presentations that are subdivided into 4 clinical types. We present a case of Waardenburg type 2 and briefly discuss the genetic basis of phenotypic expression of WS.

CASE REPORT An 18-year-old woman undergoing evaluation of intercostal neuralgia was incidentally noted to have one blue and one brown eye (figure). Both pupils reacted equally to light and accommodation. The patient's mother reported that at birth both her eyes were blue but after 2 weeks the left eye changed to brown while the right eye remained blue. At birth she had a tuft of white hair on the frontal area. She had profound sensorineural hearing loss since birth and communicated via sign language. No other family member had this condition. Her mother's eyes were brown and the color of her father's eyes was unknown. The patient was subsequently lost to follow-up.

The differential diagnosis included oculocutaneous albinism, piebaldism, oculocerebral hypopigmentation syndrome, congenital Horner syndrome, neuroblastoma, and WS. Oculocutaneous albinism is characterized by generalized hypopigmentation of skin, hair, and eyes due to defective synthesis of melanin.¹ Since the

underlying defect is in pigment production rather than absence of melanocytes, these patients do not have sensorineural hearing loss. Piebaldism is characterized by a white forelock and multiple symmetric hypopigmented or depigmented macules but without developmental anomalies of the cochlea or interocular areas.² The oculocerebral hypopigmentation syndrome is an autosomal recessive disorder characterized by the absence of pigmentation of skin and hair, microphthalmia, corneal clouding, and spastic paraplegia.³ Congenital Horner syndrome and neuroblastoma are rare causes of heterochromia irides, but no ptosis, miosis, or anhidrosis was observed in our patient. Moreover, neither of these conditions is associated with deaf-mutism or hair or skin hypopigmentation anomalies.

Given the presence of auditory and pigmentary findings since birth, a congenital syndrome involving melanocytic dysfunction was considered. The patient met 3 major and 1 minor Waardenburg Consortium diagnostic criteria⁴ (table) and a clinical diagnosis of WS type 2 was made.

DISCUSSION WS is a rare condition occurring due to maldevelopment of neural crest-derived tissues resulting in auditory, pigmentary, interocular, limb musculature, and enteric ganglia anomalies.

Clinical features. Four types of WS have been described, although phenotypic presentation may differ due to variable penetrance of the disease traits.⁴ WS1 and WS2 are characterized by one or more physical

Figure Heterochromia irides without dystopia canthorum in an 18-year-old woman



Table Waardenburg Consortium diagnostic criteria⁴

Major	Minor
Congenital sensorineural hearing loss	Congenital leukoderma, synophrys, or medial eyebrow flare
Pigmentary disturbances of the iris	Broad high nasal root
Hair hypopigmentation	Hypoplasia of alae nasi
Affected first-degree relative	Prematurely graying hair
Dystopia canthorum	

From the Mason Eye Institute (C.R.S.), and Department of Neurology (G.K.), University of Missouri-Healthcare, Columbia.

Disclosure: The authors report no disclosures.

findings of the disease (table). The distinguishing feature between the 2 types is presence or absence of dystopia canthorum—an unusual facial feature in which the medial canthi of the eyes are spaced farther apart than normal, resulting in an appearance of widely spaced eyes. Ninety-eight percent of patients with WS1 have dystopia canthorum but patients with WS2 do not have this finding.⁴ WS3 (Waardenburg-Klein) phenotypes have hypoplasia of limb musculature and/or contractures of elbows and fingers,⁵ and WS4 (Waardenburg-Shah) phenotypes have Hirschsprung disease in addition to the other common features of WS.⁶

Etiology. Melanocyte dysfunction may be due to failure of neural crest differentiation, migration, or terminal differentiation and survival in target locations. Neural crest-derived melanocytes are essential to stria vascularis of the cochlea,⁷ which explains the sensorineural hearing loss observed in WS and preserved hearing in oculocutaneous albinism. Since retinal melanocytes are not derived from the neural crest but originate from the optic cup of the developing forebrain,⁷ they may serve as an important marker for differentiation of melanocyte-specific disorders as opposed to neurocristopathies. The limb muscles, frontal bone, and enteric ganglia are derived from the neural crest, which explains the phenotypic involvement of these structures in WS1, WS3, and WS4.

WS1 and WS3 are transmitted in an autosomal dominant fashion and nearly all cases have mutations in *PAX-3* gene belonging to paired box family of transcription factors.⁷ Transcripts of *PAX-3* are expressed in cells of the neural crest between day 8 and 17 of gestation.⁷ It is hypothesized that effective levels of *PAX-3* protein determine the phenotype variation. Mild reduction in these proteins results in dystopia canthorum, 50% reduction results in melanocyte defects, and severe reduction results in limb anomalies.⁷ Hence WS1 and WS3 may actually represent phenotypic variations of the *PAX-3* mutation depending upon the severity of suppression of the transcripts of *PAX-3*. WS2 is also transmitted in an autosomal dominant fashion, and about 15% of cases have a mutation in the microphthalmia gene *MITF*, while the molecular basis of the remainder of 85% is still unknown.⁸ *MITF* appears to be a master gene controlling melanocyte differentiation, and therefore WS2 is, at least in part, melanocyte specific.

WS4 is transmitted in an autosomal recessive manner and results from homozygous mutation of endothelin-3 (*EDN3*), its receptor (*EDNRB*), or *SOX-10* (SRY-related HMG-box).⁸ Patients with heterozygous mutation of *EDN3* or *EDNRB* have isolated Hirschsprung disease.⁸ The *SOX10* gene directs the activity of other genes that signal neural crest to differentiate into melanocytes and enteric nerves.⁸ This explains the presence of Hirschsprung disease in patients with WS4.

Prognosis. Patients with WS have normal intelligence and they perform well on standardized speech tests following cochlear implantation and auditory rehabilitation. Early auditory rehabilitation is essential for normal cognitive development and is associated with excellent prognosis.⁹ Given the possibility of neural tube defects in patients with WS, folic acid supplementation is recommended in women with increased risk of having a child with WS.¹⁰

REFERENCES

1. Biswas S, Lloyd IC. Oculocutaneous albinism. *Arch Dis Child* 1999;80:565–569.
2. Thomas I, Kihiczak GG, Fox MD, Janniger CK, Schwartz RA. Piebaldism: an update. *Int J Dermatol* 2004;43:716–719.
3. Tezcan I, Demir E, Asan E, Kale G, Muftuoglu SF, Kotiloglu E. A new case of oculocerebral hypopigmentation syndrome (Cross syndrome) with additional findings. *Clin Genet* 1997;51:118–121.
4. Farrer LA, Grundfast KM, Amos J, et al. Waardenburg syndrome (WS) type I is caused by defects at multiple loci, one of which is near *ALPP* on chromosome 2: first report of the WS Consortium. *Am J Hum Genet* 1992;50:902–913.
5. Klein D. Historical background and evidence for dominant inheritance of the Klein-Waardenburg syndrome (type III). *Am J Med Genet* 1983;14:231–239.
6. Shah KN, Dalal SJ, Desai MP, Sheth PN, Joshi NC, Ambani LM. White forelock, pigmentary disorder of irides, and long segment Hirschsprung disease: possible variant of Waardenburg syndrome. *J Pediatr* 1981;99:432–435.
7. Read AP, Newton VE. Waardenburg syndrome. *J Med Genet* 1997;34:656–665.
8. Bondurand N, Dastot-Le Moal F, Stanchina L, et al. Deletions at the *SOX10* gene locus cause Waardenburg syndrome types 2 and 4. *Am J Hum Genet* 2007;81:1169–1185.
9. Cullen RD, Zdanski C, Roush P, et al. Cochlear implants in Waardenburg syndrome. *Laryngoscope* 2006;116:1273–1275.
10. Fleming A, Copp AJ. Embryonic folate metabolism and mouse neural tube defects. *Science* 1998;280:2107–2109.

Neurology[®]

Child Neurology: A patient with dissimilar eye color and deafness

Chetan R. Soni and Gyanendra Kumar

Neurology 2010;74:e25-e26

DOI 10.1212/WNL.0b013e3181d0cc5f

This information is current as of February 22, 2010

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/74/8/e25.full
References	This article cites 10 articles, 3 of which you can access for free at: http://n.neurology.org/content/74/8/e25.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/74/8/e25.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology All Neuro-ophthalmology http://n.neurology.org/cgi/collection/all_neuroophthalmology All Neurotology http://n.neurology.org/cgi/collection/all_neurotology Developmental disorders http://n.neurology.org/cgi/collection/developmental_disorders
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

