Tadpole pupil

Figure Patient's pupils

(Top) On examination, the pupils are round, isocoric, and reactive to light. (Bottom) On this self-taken photograph, the left pupil is no longer round and is larger than the right pupil. Focal dilation of the superonasal sector of the left pupil is readily visible.

A young woman has recurrent episodes of unilateral mydriasis lasting several minutes. The left pupil becomes large and deformed, described as “egg-shaped” (figure). Examination, including pharmacologic pupil tests, revealed no ophthalmologic or neurologic abnormalities. Cranio-orbital MRI was normal. Diagnosis of tadpole pupil was confirmed by self-taken photographs using a cellular phone.

Spontaneous segmental spasm of the iris dilator can cause pupillary distortion that resembles the shape of a tadpole.^{1,2} Occasional patients with tadpole pupil have underlying Horner syndrome; otherwise, the syndrome is benign and self-limited.¹ Cellular phone “telemedicine” was particularly helpful in diagnosing this form of episodic mydriasis.

Aki Kawasaki, MD, Cedric Mayer, MD, Lausanne, Switzerland

A. Kawasaki has received honoraria from serving on a scientific advisory panel for Bayer SpA. C. Mayer reports no disclosures. **Go to Neurology.org for full disclosures.**

Correspondence & reprint requests to Dr. Kawasaki: aki.kawasaki@fa2.ch

1. Thompson HS, Zackon DH, Czarnecki JS. Tadpole-shaped pupils caused by segmental spasm of the iris dilator muscle. *Am J Ophthalmol* 1983;96:467–477.
2. Balaggan KS, Hugkulstone CE, Bremner FD. Episodic segmental iris dilator muscle spasm: the tadpole-shaped pupil. *Arch Ophthalmol* 2003;121:744–745.