


Teaching NeuroImages:

Diagnostic utility of FDG-PET in neurolymphomatosis

Michel Toledano, MD
Mustaqem A. Siddiqui,
MD, MBA
Carrie A. Thompson, MD
Ivan Garza, MD
Sean J. Pittock, MD

Correspondence to
Dr. Toledano:
toledano.michel@mayo.edu

Figure Face, orbit, and neck imaging



Axial MRI T1 gadolinium images without any contrast enhancement at the level of the trigeminal nerve (A) or parotid gland (D). PET reveals a focus of intense fluorodeoxyglucose (FDG) avidity extending through the foramen ovale along the course of the left V3 branch of the trigeminal nerve in both axial (B) and coronal (C) cuts, consistent with perineural spread of disease. Axial (E) and coronal (F) cuts show increased FDG activity in the left parotid gland.

A 70-year-old woman with a history of non-Hodgkin lymphoma presented with left-sided facial pain. MRI of the face, orbit, and neck was negative (figure). A week later, she developed hypesthesia in a V3 distribution. Repeat MRI was again non-diagnostic but fluorodeoxyglucose (FDG)-PET showed increased uptake along the left V3 branch of the trigeminal nerve, as well as in the parotid gland. Biopsy of the left parotid confirmed recurrent lymphoma.

Neurolymphomatosis is a rare manifestation of hematologic disease and diagnosis is often delayed.¹ Our case illustrates the utility of FDG-PET in establishing the diagnosis of neurolymphomatosis when suspicion is high but MRI is unrevealing.²

AUTHOR CONTRIBUTIONS

Drs. Toledano, Garza, and Pittock made substantive contributions to the design of the study and drafting of the manuscript. Drs. Siddiqui and Thompson made substantive contributions to the revision of the manuscript.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

REFERENCES

1. Grisariu S, Avni B, Batchelor TT, et al. Neurolymphomatosis: an International Primary CNS lymphoma Collaborative Group report. *Blood* 2010;115:5005–5011.
2. Salm LP, Van der Hiel B, Stokkel MP. Increasing importance of 18F-FDG PET in the diagnosis of neurolymphomatosis. *Nucl Med Commun* 2012;33:907–916.

Download teaching slides:
www.neurology.org

From the Department of Neurology (M.T., I.G., S.J.P.), Division of Hospital Internal Medicine, Internal Medicine (M.A.S.), and Department of Hematology (C.A.T.), Mayo Clinic, Rochester, MN.

Neurology®

Teaching *NeuroImages*: Diagnostic utility of FDG-PET in neurolymphomatosis

Michel Toledano, Mustaqeem A. Siddiqui, Carrie A. Thompson, et al.

Neurology 2013;81:e3

DOI 10.1212/WNL.0b013e318297eef5

This information is current as of July 1, 2013

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/81/1/e3.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2013/06/29/81.1.e3.DC1
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/81/1/e3.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Hematologic PET http://n.neurology.org/cgi/collection/hematologic PET http://n.neurology.org/cgi/collection/pet
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

