

**Editors' Note:** The 1994 guidelines for the diagnosis of the vegetative state (VS) should be revised, Young and Owen suggest. Estraneo et al. agree, pointing to the development of new, advanced tools that improve recognition of discernible intentional responses in patients with prolonged anoxic VS. Some corrections and interesting additions to the historical article on cerebral PET scanning are made by Singhal. Okun et al. agree and explain the challenge of word count limit.

*Chafic Karam, MD, and Robert C. Griggs, MD*

#### PREDICTORS OF RECOVERY OF RESPONSIVENESS IN PROLONGED ANOXIC VEGETATIVE STATE

**G. Bryan Young, Adrian M. Owen, London, Canada:**

Estraneo et al.<sup>1</sup> prospectively studied 43 patients post cardiac arrest who had been clinically vegetative for over a month. Nine (21%) of the patients regained behavioral responses—although still disabled in motor function—when the period of observation was extended to 23–26 months.

With the use of advanced technology—fMRI, event-related potentials, and quantitative EEG measures of responsiveness<sup>2–4</sup>—the yield of responsive patients may have been higher. The additional yield of such responses is approximately 17% in patients deemed vegetative for much longer.<sup>3</sup> It is possible that another 6 patients of the 34 behaviorally unresponsive patients in this study<sup>1</sup> might have shown responses on further testing.

Although patients may have severe motor disabilities, it is important to be able to communicate with patients to determine their needs and wishes and for families to know that their injured loved ones are aware. Thus, precision in diagnosis and prognosis is vital.

The 1994 guidelines<sup>5</sup> for the diagnosis of VS should be revised. Patients require much longer follow-ups and, we argue, the use of state-of-the-art brain imaging techniques before a firm diagnosis of VS is made.

**Author Response: Anna Estraneo, Pasquale Moretta, Telesse Terme; Luigi Trojano, Caserta, Italy:** Detecting signs of covert cognition in VS is difficult

for clinicians. Technologically advanced tools have been developed to improve recognition of discernible, intentional (nonreflexive) responses in such patients.<sup>6</sup> We agree that the use of more sensitive methods could help identify a higher number of responsive patients and to redefine classical diagnostic criteria. This seems to be particularly relevant since it appears that clinical evolution of VS is changing, and that, for instance, “late recovery” of responsiveness and consciousness can no longer be regarded as exceptional.<sup>7</sup> However, such technological advances present several limitations: only a selected sample of patients’ clinical features can be assessed; study paradigms and methods of analysis are complex; and acquisition and management costs are high.<sup>8,9</sup> These restrictions prohibit large-scale studies aimed to assess diagnostic sensitivity and specificity of modern technologies. Widely available diagnostic tools should be studied to assess their possible prognostic value and to provide reliable information for clinical decision-making and treatment management.<sup>1</sup>

© 2013 American Academy of Neurology

1. Estraneo A, Moretta P, Loreto V, et al. Predictors of recovery of responsiveness in prolonged anoxic vegetative state. *Neurology* 2013;80:464–470.
2. Naci L, Monti MM, Cruse D, et al. Brain-computer interfaces for communication with nonresponsive patients. *Ann Neurol* 2012;72:312–323.
3. Monti MM, Vanhaudenhuyse A, Coleman MR, et al. Willful modulation of brain activity and communication in disorders of consciousness. *N Engl J Med* 2010;362:579–589.
4. Cruse D, Chennu S, Chatelle C, et al. Bedside detection of awareness in the vegetative state: a cohort study. *Lancet* 2011;378:2088–2094.
5. The Multi-Society Task force on PVS. Medical aspects of the vegetative state: The Multi-Society Task Force on PVS. *N Engl J Med* 1994;330:1499–1508.
6. Owen AM, Schiff ND, Laureys S. A new era of coma and consciousness science. *Prog Brain Res* 2009;177:399–411.
7. Bekinschtein TA, Manes FF, Villarreal M, et al. Functional imaging reveals movement preparatory activity in the vegetative state. *Front Hum Neurosci* 2011;27:1–11.
8. Bardin JC, Fins JJ, Katz DI, et al. Dissociations between behavioural and functional magnetic resonance imaging-based evaluations of cognitive function after brain injury. *Brain* 2011;134:769–782.

9. Estraneo A, Moretta P, Loreto V, et al. Late recovery after traumatic, anoxic, or hemorrhagic long-lasting vegetative state. *Neurology* 2010;75:239–245.

### THE HISTORY OF CEREBRAL PET SCANNING: FROM PHYSIOLOGY TO CUTTING-EDGE TECHNOLOGY

**Tarun Singhal, Boston:** The title of the article by Portnow et al.<sup>1</sup> is not suitable because there is uneven coverage of the development of PET imaging devices and inadequate coverage of radiopharmaceuticals or their translational applications. There are some key conceptual and factual errors as well.

Contrary to the authors' contention, half-life of Carbon-14 is approximately 5,730 years, while half-life of Fluorine-18 is only 110 minutes: the latter's half-life is shorter—not longer—than the former's. Moreover, there was a need for an alternative agent for glucose imaging because Carbon-14 decays by beta particle formation and beta particles cannot penetrate the human body for image formation. Gamma rays formed after the positron decay of fluorine-18 and other positron emitters can penetrate the human body to enable emission imaging in living humans. Additionally, authors show an image of a <sup>18</sup>F-spiperone scan but do not mention the pioneering studies by Wagner et al.<sup>2,3</sup> on neuroreceptor imaging in the brain with 3-N-[<sup>11</sup>C] methylspiperone. Dr. Wagner is considered a forefather of nuclear medicine.<sup>4</sup>

Finally, PET can also be used for cerebellar and brain-stem imaging—in addition to cerebral imaging—which is of relevance for several brain disorders.

**Author Response: Michael Okun, Leah Portnow, David Vaillancourt, Gainesville, FL:** We appreciate the comments by Dr. Singhal. Dr. Singhal is correct that we should have used the word “shorter half-life” instead of “longer half-life” when describing the half-life of FDG compared with <sup>14</sup>CDG. We are grateful that this error was noticed. We also agree that other elements of PET imaging including radiopharmaceuticals and the translational applications should have been included. In prior drafts, we had a more developed version consistent with these suggestions, but with the word count limits we were constrained to focus on key areas that were of particular interest to our research.

**Editors' Note:** A correction regarding the half-life of FDG appears on page 1275.

© 2013 American Academy of Neurology

1. Portnow LH, Vaillancourt DE, Okun MS. The history of cerebral PET scanning: from physiology to cutting-edge technology. *Neurology* 2013;80:952–956.
2. Wagner HN Jr, Burns HD, Dannals RF, et al. Imaging dopamine receptors in the human brain by positron tomography. *Science* 1983;221:1264–1266.
3. Jones T, Rabiner EA. The development, past achievements, and future directions of brain PET. *J Cereb Blood Flow Metab* 2012;32:1426–1454.
4. Society of Nuclear Medicine and Molecular Imaging member news, September 27, 2012. Available at: <http://www.snm.org/index.cfm?PageID=12082>. Accessed July 31, 2013.

### CORRECTION

#### The history of cerebral PET scanning: From physiology to cutting-edge technology

In the Historical Neurology article “The history of cerebral PET scanning: From physiology to cutting-edge technology” by L.H. Portnow et al. (*Neurology*® 2013;80:952–956), there is an error on page 954. When describing the half-life of FDG compared with [<sup>14</sup>C]DG, the authors should have used “shorter half-life” instead of “longer half-life.” The authors regret the error.

Author disclosures are available upon request ([journal@neurology.org](mailto:journal@neurology.org)).

# Neurology®

## **Predictors of recovery of responsiveness in prolonged anoxic vegetative state**

G. Bryan Young, Anna Estraneo, Adrian M. Owen, et al.

*Neurology* 2013;81;1274-1275

DOI 10.1212/WNL.0b013e3182a7ae28

**This information is current as of September 30, 2013**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://n.neurology.org/content/81/14/1274.full">http://n.neurology.org/content/81/14/1274.full</a>
<b>References</b>	This article cites 9 articles, 2 of which you can access for free at: <a href="http://n.neurology.org/content/81/14/1274.full#ref-list-1">http://n.neurology.org/content/81/14/1274.full#ref-list-1</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.neurology.org/about/about_the_journal#permissions">http://www.neurology.org/about/about_the_journal#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://n.neurology.org/subscribers/advertise">http://n.neurology.org/subscribers/advertise</a>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

