


Teaching NeuroImages: Concurrent hemifacial spasm and trigeminal neuralgia due to vertebrobasilar dolichoectasia

Shaheen E. Lakhan, MD,
PhD, MS, Med


Correspondence to
Dr. Lakhan:
slakhan@gnif.org

Figure 1 Brain MRI, 3D magnetization-prepared rapid acquisition, axial view



MRI demonstrates dolichoectatic basilar artery distorting the right trigeminal nerve at the level of the mid pons (arrow). For reference, contralateral trigeminal nerve displayed with asterisk.

Figure 2 Brain MRI, 3D magnetization-prepared rapid acquisition, axial view



MRI demonstrates dolichoectatic basilar artery distorting the right facial nerve at the cerebellopontine angle (arrow). For reference, contralateral facial nerve displayed with asterisk.

A 64-year-old man reported a 2-week worsening of a 2-year history of paroxysmal, lancinating right facial pain and spasms. Head CT demonstrated a tortuous and ectatic vertebral-basilar system. MRI confirmed a dolichoectatic basilar artery that distorted the right trigeminal (figure 1) and facial nerve roots (figure 2). He failed conservative medical management including carbamazepine, oxcarbazepine, and levetiracetam. He underwent Gamma Knife radiosurgery with a few months of pain relief and was offered botulinum neurotoxin injection for spasms.

This case represents a rare presentation of concurrent hemifacial spasm and trigeminal neuralgia linked to an ectatic basilar artery.¹

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

Dr. Lakhan serves on the American Academy of Neurology's Distance Learning Subcommittee and is an editorial team member of the Resident & Fellow Section of *Neurology*[®]. Go to Neurology.org for full disclosures.

REFERENCE

1. Kwon HM, Lee YS. Dolichoectasia of the intracranial arteries. *Curr Treat Options Cardiovasc Med* 2011;13:261–267.

Neurology®

Teaching NeuroImages: Concurrent hemifacial spasm and trigeminal neuralgia due to vertebrobasilar dolichoectasia

Shaheen E. Lakhani

Neurology 2013;81:e52

DOI 10.1212/WNL.0b013e3182a1aeb

This information is current as of August 19, 2013

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/81/8/e52.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2013/08/14/81.8.e52.DC1
References	This article cites 1 articles, 0 of which you can access for free at: http://n.neurology.org/content/81/8/e52.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Movement Disorders http://n.neurology.org/cgi/collection/all_movement_disorders Other cerebrovascular disease/ Stroke http://n.neurology.org/cgi/collection/other_cerebrovascular_disease__stroke Trigeminal neuralgia http://n.neurology.org/cgi/collection/trigeminal_neuralgia
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

