

acid sequence “HRGMLR” was analyzed by the BLAST program by comparing the sequences of moesin and CMV proteins in our study, was located at a different site from the C-terminal, and was listed as the candidate of immune target. Separately, we agree that an animal model of AIDP by immunization with moesin protein (peptide) is necessary to confirm the disease pathogenesis of CMV-related AIDP, and intend to do so in the future.

© 2014 American Academy of Neurology

1. Sawai S, Satoh M, Mori M, et al. Moesin is a possible target molecule for cytomegalovirus-related Guillain-Barré syndrome. *Neurology* 2014;83:113–117.
2. Miyaji K, Shahrizaila N, Umaphathi T, et al. Are ERM (ezrin/radixin/moesin) proteins targets for autoantibodies in demyelinating neuropathies? *Hum Immunol* 2014;75:1089–1091.
3. Hafer-Macko CE, Sheikh KA, Li CY, et al. Immune attack on the Schwann cell surface in acute inflammatory demyelinating polyneuropathy. *Ann Neurol* 1996;39:625–635.
4. Tsukita S, Yonemura S. Cortical actin organization: lessons from ERM (ezrin/radixin/moesin) proteins. *J Biol Chem* 1999;274:34507–34510.
5. Amar S, Oyaisu K, Li L, et al. Moesin: a potential LPS receptor on human monocytes. *J Endotoxin Res* 2001;7:281–286.
6. Iontcheva I, Amar S, Zawawi KH, et al. Role for moesin in lipopolysaccharide-stimulated signal transduction. *Infect Immun* 2004;72:2312–2320.
7. Takamatsu H, Espinoza JL, Lu X, et al. Anti-moesin antibodies in the serum of patients with aplastic anemia stimulate peripheral blood mononuclear cells to secrete TNF-alpha and IFN-gamma. *J Immunol* 2009;182:703–710.
8. Takamatsu H, Feng X, Chuhjo T, et al. Specific antibodies to moesin, a membrane-cytoskeleton linker protein, are frequently detected in patients with acquired aplastic anemia. *Blood* 2007;109:2514–2520.
9. Yuki N, Taki T, Inagaki F, et al. A bacterium lipopolysaccharide that elicits Guillain-Barré syndrome has a GM1 ganglioside-like structure. *J Exp Med* 1993;178:1771–1775.
10. Yuki N, Yamada M, Koga M, et al. Animal model of axonal Guillain-Barré syndrome induced by sensitization with GM1 ganglioside. *Ann Neurol* 2001;49:712–720.
11. Goodfellow JA, Bowes T, Sheikh K, et al. Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy. *J Neurosci* 2005;25:1620–1628.
12. Greenshields KN, Halstead SK, Zitman FMP, et al. The neuropathic potential of anti-GM1 autoantibodies is regulated by the local glycolipid environment in mice. *J Clin Invest* 2009;119:595–610.

CORRECTIONS

Occupational solvent exposure and cognition: Does the association vary by level of education?

In the article “Occupational solvent exposure and cognition: Does the association vary by level of education?” by E.L. Sabbath et al. (*Neurology*® 2012;78:1754–1760), there is an error in the Methods. The authors listed hydrazine as an example of petroleum solvent and toluene diisocyanate as an example of a nonbenzene aromatic solvent; however, toluene diisocyanate and hydrazine were not included in any analyses because they are not solvents. The first sentence under “Exposures” should have read: “A validated job exposure matrix (JEM) was used to characterize exposure to 4 solvent categories: chlorinated solvents (all species combined, plus separate assessments of tetrachloromethane, trichloroethylene, perchloroethylene, dichloromethane, trichloroethane), petroleum solvents (all species combined), benzene, and nonbenzene aromatic solvents (all species combined).” The results and conclusions of the paper are not affected by this correction. The authors regret the error.

Time may not fully attenuate solvent-associated cognitive deficits in highly exposed workers

In the article “Time may not fully attenuate solvent-associated cognitive deficits in highly exposed workers” by E.L. Sabbath et al. (*Neurology*® 2014;82:1716–1723), there is an error in the Methods. The authors listed hydrazine as an example of petroleum solvent and toluene diisocyanate as an example of a nonbenzene aromatic solvent; however, toluene diisocyanate and hydrazine were not included in any analyses because they are not solvents. The first sentence under “Exposure” should have read: “A job exposure matrix (JEM) was used to characterize lifetime exposure to 4 categories of organic solvents: chlorinated solvents (all species combined, plus separate assessments of tetrachloromethane, trichloroethylene, perchloroethylene, dichloromethane, trichloroethane), petroleum solvents (all species combined), benzene, and nonbenzene aromatic solvents (all species combined).” The results and conclusions of the paper are not affected by this correction. The authors regret the error.

Neurology®

Time may not fully attenuate solvent-associated cognitive deficits in highly exposed workers

Neurology 2014;83:2315
DOI 10.1212/WNL.0000000000001103

This information is current as of December 8, 2014

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/83/24/2315.2.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

