Increased CSF biomarkers of angiogenesis in Parkinson disease

OPEN

Shorena Janelidze, PhD*
 Daniel Lindqvist, MD,
 PhD*
 Veronica Francardo, PhD
 Sara Hall, MD
 Henrik Zetterberg, MD,
 PhD
 Kaj Blennow, MD, PhD
 Charles H. Adler, MD,
 PhD
 Thomas G. Beach, MD,
 PhD
 Geidy E. Serrano, PhD
 Danielle van Westen,
 MD, PhD
 Elisabet Londos, MD,
 PhD
 M. Angela Cenci, MD,
 PhD
 Oskar Hansson, MD,
 PhD

Correspondence to
 Dr. Lindqvist:
 Daniel.Lindqvist@med.lu.se

ABSTRACT

Objective: To study biomarkers of angiogenesis in Parkinson disease (PD), and how these are associated with clinical characteristics, blood-brain barrier (BBB) permeability, and cerebrovascular disease.

Methods: In this cross-sectional analysis, 38 elderly controls and 100 patients with PD (82 without dementia and 18 with dementia) were included from the prospective Swedish BioFinder study. CSF samples were analyzed for the angiogenesis biomarkers vascular endothelial growth factor (VEGF); its receptors, VEGFR-1 and VEGFR-2; placental growth factor (PlGF); angiopoietin 2 (Ang2); and interleukin-8. BBB permeability, white matter lesions (WMLs), and cerebral microbleeds (CMB) were assessed. CSF angiogenesis biomarkers were also measured in 2 validation cohorts: (1) 64 controls and 87 patients with PD with dementia; and (2) 35 controls and 93 patients with neuropathologically confirmed diagnosis of PD with and without dementia.

Results: Patients with PD without dementia displayed higher CSF levels of VEGF, PlGF, and sVEGFR-2, and lower levels of Ang2, compared to controls. Similar alterations in VEGF, PlGF, and Ang2 levels were observed in patients with PD with dementia. Angiogenesis markers were associated with gait difficulties and orthostatic hypotension as well as with more pronounced BBB permeability, WMLs, and CMB. Moreover, higher levels of VEGF and PlGF levels were associated with increased CSF levels of neurofilament light (a marker of neurodegeneration) and monocyte chemoattractant protein-1 (a marker of glial activation). The main results were validated in the 2 additional cohorts.

Conclusions: CSF biomarkers of angiogenesis are increased in PD, and they are associated with gait difficulties, BBB dysfunction, WMLs, and CMB. Abnormal angiogenesis may be important in PD pathogenesis and contribute to dopa-resistant symptoms. *Neurology*® 2015;85:1834-1842

GLOSSARY

Aβ = β-amyloid; **Ang** = angiopoietin; **ARWMC** = age-related white matter changes; **BBB** = blood-brain barrier; **CMB** = cerebral microbleeds; **CV** = coefficient of variation; **DLB** = dementia with Lewy bodies; **IL** = interleukin; **MCP-1** = monocyte chemoattractant protein-1; **NFL** = neurofilament light; **p-tau** = phospho-tau; **PD** = Parkinson disease; **PDD** = Parkinson disease with dementia; **PDND** = Parkinson disease without dementia; **PlGF** = placental growth factor; **SNpc** = substantia nigra pars compacta; **t-tau** = total tau; **VEGF** = vascular endothelial growth factor; **WML** = white matter lesions.

Angiogenesis might be an important mechanism involved in pathophysiology of Parkinson disease (PD). Increased numbers of endothelial cell nuclei and blood vessels have been found postmortem in the substantia nigra pars compacta (SNpc) of patients with PD^{1,2} and parkinsonian primates.³ PD status at the time of death is also associated with greater expression of integrin αvβ3, a selective marker of angiogenic endothelial cells, in the SNpc, the locus ceruleus, and putamen.⁴

*These authors contributed equally to this work.

From the Clinical Memory Research Unit, Department of Clinical Sciences (S.J., E.L., O.H.), Lund University, Malmö; the Department of Clinical Sciences (D.L., S.H.), Division of Psychiatry (D.L.), Department of Experimental Medical Science (V.F., M.A.C.), and Clinical Sciences, Diagnostic Radiology (D.v.W.), Lund University, Lund; Psychiatry Skåne (D.L.), Lund; the Department of Neurology (S.H.) and Memory Clinic (E.L., O.H.), Skåne University Hospital, Lund; Institute of Neuroscience and Physiology (H.Z., K.B.), Department of Psychiatry and Neurochemistry, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden; UCL Institute of Neurology (H.Z.), London, UK; The Torsten Söderberg Professorship in Medicine at the Royal Swedish Academy of Sciences (K.B.), Stockholm, Sweden; Department of Neurology (C.H.A.), Mayo Clinic, Scottsdale; Banner Sun Health Research Institute (T.G.B., G.E.S.), Sun City, AZ; and Imaging and Function (D.v.W.), Skåne University Health Care, Lund, Sweden.

Go to Neurology.org for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article. The Article Processing Charge was paid by Lund University, Sweden, with funding from the European Research Council.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Editorial, page 1826

Supplemental data
 at Neurology.org

Angiogenesis is regulated by a plethora of different proteins, including members of the vascular endothelial growth factor (VEGF) family (e.g., VEGF and its receptors, VEGFR-1 and VEGFR-2, placental growth factor [PlGF]),⁵ the angiopoietin (Ang) family (e.g., Ang2),⁶ and proinflammatory chemokines (e.g., interleukin [IL]–8).⁷

Previous postmortem studies exploring the role of angiogenesis in PD have included few cases.^{1,2} Consequently, there is a need to determine whether biomarkers of angiogenesis are reliably changed in PD. Moreover, the contribution of aberrant angiogenesis to specific aspects of PD symptomatology (such as cognitive or motor dysfunction) has not been investigated. It would be also interesting to establish if changes in angiogenesis biomarkers are associated with overall cerebrovascular pathology. To this end, we measured several biomarkers of angiogenesis in CSF samples from a large cohort of patients with PD with and without dementia and healthy controls. We investigated cross-sectional associations between the measured angiogenesis markers and disease symptoms, blood–brain barrier (BBB) permeability, white matter lesions (WML), and cerebral microbleeds (CMB). Finally, the main findings of the study were replicated in 2 additional cohorts of healthy controls and patients with PD, of which one consisted of neuropathologically confirmed cases.

METHODS **Standard protocol approvals, registrations, and patient consents.** The Ethics Committee of Lund University approved both studies originating from Lund. Study participants gave informed consent to research. The study was conducted in accordance with the provisions of the Helsinki Declaration.

Study population. **Cohort 1.** One hundred patients with PD and 38 healthy controls were enrolled in the prospective Swedish BioFinder study (www.biofinder.se) at the Neurology Clinic, Skåne University Hospital, Lund, Sweden. The PD group included 82 patients without dementia (PDND) and 18 patients with dementia (PDD). PD diagnosis was set according to the National Institute of Neurological Disorders and Stroke Diagnostic Criteria.⁸ PDD was diagnosed according to the Clinical Diagnostic Criteria for Dementia Associated with Parkinson's Disease.⁹

Analysis of the data from cohort 1 indicated that to provide 90% power at an α level of 0.05, a total of 100 patients would be required.

Cohort 2. The first validation cohort included 64 healthy controls and 87 patients (25 PDD and 62 dementia with Lewy bodies [DLB]), all assessed at the Memory Clinic, Skåne University Hospital, Lund, Sweden.

Cohort 3. The second independent validation cohort included 35 healthy controls, 27 PDND patients, and 66 PDD

patients, all neuropathologically confirmed cases selected by the Arizona Parkinson's Disease Consortium National Brain and Tissue Resource for Parkinson's Disease and Related Disorders. All participants or their legal representatives signed an Institutional Review Board–approved informed consent form before the time of death. The PDD group included 32 patients without AD-related histopathology and 34 patients with sufficient load of plaques and tangles to meet criteria for AD diagnosis. The 2 dementia groups were finally merged into one single group of PDD cases because the concurrent AD pathology was found not to have any impact on the parameters under investigation.

Detailed information on all the study cohorts is available in the supplemental data on the *Neurology*[®] Web site at Neurology.org.

CSF sampling and biological assays. In cohort 1 and cohort 2, CSF was collected as described previously.¹⁰ In cohort 3 with neuropathology confirmed cases, postmortem CSF was collected from the lateral ventricles while the brain was still in situ, after removal of the skullcap. VEGF, PlGF, sVEGFR-1, sVEGFR-2, Ang2, and IL-8 were quantified in CSF samples from cohort 1 using multiplex electrochemiluminescence immunoassay (Meso Scale Discovery, Gaithersburg, MD; supplemental data). All the samples were measured in duplicate and the mean of the duplicate was used in the statistical analysis. Detection limits are provided in the supplemental data. The coefficient of variation (CV) was below 20% for all assays. The few samples with CV >20% did not affect the results and were therefore included in the statistical analysis. In the validation cohort, VEGF, PlGF, and sVEGFR-1 were measured using the same methodology as above. For further validation purposes, CSF levels of VEGF were also measured using VEGF V-PLEX kit using other antibodies than the VEGF assay mentioned above (see supplemental data for details).

The albumin ratio was calculated as CSF albumin (mg/L)/plasma albumin (g/L) and was used as a measure of the BBB function. Albumin and monocyte chemoattractant protein-1 (MCP-1) were measured as described previously.^{11,12} Total tau (t-tau; ADx Neurosciences NV, Gent, Belgium), neurofilament light (NFL; Uman Diagnostics, Umeå, Sweden), phospho-tau (p-tau), and β -amyloid (A β) 42 (Fujirebio Europe, Gent, Belgium) were analyzed using commercially available ELISA kits.

MRI. In cohort 1, 32 controls and 58 patients with PD underwent MRI scans using a 3T Siemens[®] system to quantify WMLs and CMB (see supplemental data for details).

Statistical analyses. The Statistical Package for the Social Sciences (SPSS Inc., Chicago, IL) was used for statistical calculations. Due to skewness, all CSF biomarkers were transformed into their natural logarithm before analyses. Untransformed and unadjusted values are shown in table 1, tables e-1 and e-2. For groupwise comparisons, we used Student *t* test or one-way analysis of variance adjusting for covariates. For associations between 2 continuous variables, Pearson partial correlation was used. Standard linear regressions were used to investigate associations between a continuous dependent variable and continuous or categorical independent variables.

In addition to CSF concentrations of angiogenesis biomarkers, we also used the VEGF/sVEGFR-1 and PlGF/sVEGFR-1 ratios when investigating associations with clinical data. The rationale for this is that sVEGFR-1 has direct antagonistic effect on VEGF and PlGF by sequestering the ligands from the membrane receptors.^{13–15} Consequently, high VEGF/sVEGFR-1 and PlGF/sVEGFR-1 ratios provide an index of the bioactive levels of VEGF and PlGF.

Table 1 Demographic characteristics and CSF levels of angiogenesis biomarkers in healthy controls, patients with PDND, and patients with PDD included in cohort 1

	Controls (n = 38)	PDND (n = 82)	PDD (n = 18)
Sex, F/M, n (%)	22 (58)/16	30 (37)/52 ^a	5 (28)/13 ^b
Age, y, mean ± SD	65.4 ± 8.6	64.5 ± 10.5	72.3 ± 6.0 ^c
Illness duration, y, mean ± SD	NA	6.5 ± 5.4	16.3 ± 6.4 ^d
Hoehn & Yahr, mean ± SD	NA	2.0 ± 0.8	3.1 ± 0.8 ^d
Schwab & England	100 (100-100)	90 (88-96) ^a	70 (60-80) ^c
UPDRS-III motor score	0 (0-2)	18 (11-25) ^a	36 (24-49) ^c
Tandem gait test	0 (0-0)	0 (0-1)	2 (1-3)
MMSE score	29.0 (27.0-29.3)	29.0 (28.0-29.0)	25.0 (20.5-26.3) ^c
Orthostatic hypotension, %	27.3	48.7 ^a	76.5 ^c
Cardiovascular disease, %	31.6	17.1	22.2
Diabetes mellitus, %	2.6	3.7	0.0
Asthma/COPD, %	7.9	12.2	5.6
Anti-inflammatory drugs, %	7.9	3.7	5.6
CSF plasma albumin ratio	3.5 (2.8-4.3)	4.8 (3.4-6.1) ^a	5.5 (3.9-6.0) ^b
WMLs, total ARWMC score	2.5 (0.0-5.0)	4.0 (0.0-7.0)	4.5 (2.5-7.0)
VEGF	67.9 (52.2-78.2)	75.0 (62.0-95.9) ^a	94.1 (80.7-112.9) ^b
PlGF	57.1 (46.0-69.6)	65.7 (54.2-86.0) ^a	85.9 (74.3-103.0) ^b
sVEGFR-1	118.9 (93.8-159.8)	114.8 (93.1-143.5)	106.3 (62.5-147.5)
sVEGFR-2	1.7 (1.4-2.3)	2.1 (1.9-2.7) ^a	2.4 (1.9-2.6)
VEGF/sVEGFR-1 ratio	0.6 (0.4-0.7)	0.7 (0.5-0.9)	0.9 (0.7-1.3) ^b
PlGF/sVEGFR-1 ratio	0.5 (0.4-0.7)	0.6 (0.4-0.9)	0.8 (0.7-1.4) ^c
Ang2	260.3 (197.2-350.1)	227.7 (177.3-278.9) ^a	200.9 (159.8-346.2) ^b
IL-8	30.1 (26.4-33.8)	33.9 (27.7-38.9)	38.9 (33.1-50.3) ^b

Abbreviations: Ang2 = angiopoietin 2; ARWMC = age-related white matter changes; COPD = chronic obstructive pulmonary disease; IL-8 = interleukin-8; MMSE = Mini-Mental State Examination; NA = not available; PDD = Parkinson disease with dementia; PDND = Parkinson disease without dementia; PlGF = placental growth factor; UPDRS = Unified Parkinson's Disease Rating Scale; VEGF = vascular endothelial growth factor; WMLs = white matter lesions.

Concentrations are given in pg/mL. Angiogenesis biomarkers were skewed and therefore ln-transformed before statistical analyses. Raw values of CSF biomarkers and clinical variables are presented since they are more meaningful than transformed values. Data are shown as median (interquartile range) unless otherwise specified. The *p* values are adjusted for age and sex as described in Methods when comparing angiogenesis biomarkers between groups.

^aPatients with PDND differed significantly from healthy controls (*p* < 0.05).

^bPatients with PDD differed significantly from healthy controls (*p* < 0.05) but not from patients with PDND.

^cPatients with PDD differed significantly from both patients with PDND and healthy controls (*p* < 0.05).

^dPatients with PDD differed from patients with PDND (*p* < 0.05).

In the control and patient groups, most of the analytes were positively associated with age (*p* < 0.05). There were also sex differences for VEGF, PlGF, and CSF/plasma albumin ratio (higher levels in men, *p* < 0.05). Therefore all the subsequent statistical analyses were controlled for age, whereas analyses of VEGF, PlGF, VEGF/sVEGFR-1 ratio, PlGF/sVEGFR-1 ratio, and CSF/plasma albumin ratio were also controlled for sex. In the PD group, status as de novo PD, disease duration, or levodopa equivalent were not associated with any of the measured CSF biomarkers, with the exception of IL-8, which correlated with levodopa equivalent (*p* < 0.05).

RESULTS Demographic and clinical characteristics of the study participants as well as CSF concentrations of angiogenesis biomarkers for all the cohorts are given in table 1, tables e-1 and e-2.

Cohort 1: Group comparisons. Univariate comparisons among the control, PDND, and PDD groups revealed altered levels of VEGF (*p* = 0.022), PlGF (*p* = 0.011), VEGF/sVEGFR-1 ratio (*p* = 0.013), PlGF/sVEGFR-1 ratio (*p* = 0.012), sVEGFR-2 (*p* = 0.001), Ang2 (*p* = 0.049), and IL-8 (*p* = 0.033). PDND patients displayed higher levels of VEGF (*p* = 0.012), PlGF (*p* = 0.020), and sVEGFR-2 (*p* < 0.001), and lower levels of Ang2 (*p* = 0.044), when compared to healthy controls (figure 1, A–D). Also in the PDD group, VEGF (*p* = 0.028) and PlGF (*p* = 0.005) were increased and Ang2 was decreased (*p* = 0.030) compared to controls (figure 1, A and B). Further, VEGF/sVEGFR-1 ratio (*p* = 0.004),

Figure 1 Angiogenesis biomarkers in CSF of healthy controls, PDND, and PDD patients (cohort 1)

Vascular endothelial growth factor (VEGF) (A), placental growth factor (PIGF) (B), sVEGFR-2 (C), angiopoietin 2 (Ang2) (D), VEGF/sVEGFR-1 ratio (E), and PIGF/sVEGFR-1 ratio (F) are altered in patients with Parkinson disease without dementia (PDND) and patients with Parkinson disease with dementia (PDD) compared to controls. Angiogenesis biomarkers were skewed and therefore ln-transformed before statistical analyses; p values are from analysis of covariance adjusting for age and sex as described in the Methods; data are presented as mean $\pm$ 95% confidence interval.

PIGF/sVEGFR-1 ratio ($p = 0.003$), and IL-8 ($p = 0.01$) were selectively increased in PDD cases compared to controls (figure 1, E and F). Similar results were obtained with the VEGF V-PLEX kit (supplemental data, table e-3).

Cohort 1: Angiogenesis biomarkers and clinical assessments.

In patients with PD (all participants), worse performance on the tandem gait test (indicating postural instability and gait difficulties) was associated with increased CSF levels of VEGF ($\beta = 0.22$, $p = 0.037$), PIGF ($\beta = 0.34$, $p = 0.002$), and IL-8 ($\beta = 0.23$, $p = 0.014$). We did not find any associations between angiogenesis biomarkers and cognitive performance assessed using Mini-Mental State Examination and Alzheimer's Disease Assessment Scale items 1–3. The associations between clinical symptoms and VEGF as well as PIGF were similar in the PDND group. However, in this group we also observed positive correlations between UPDRS-II item 15 (gait, off-state) and CSF VEGF ($\beta = 0.46$, $p = 0.003$) and PIGF ($\beta = 0.38$, $p = 0.022$) as well as a negative correlation between CSF Ang2 and UPDRS-III ($\beta = -0.25$, $p = 0.029$).

Orthostatic blood pressure might result in transient brain ischemia.^{16,17} PDND patients with a significant drop in diastolic blood pressure when standing had higher levels of VEGF/sVEGFR-1 ratio and PIGF/sVEGFR1 ratio compared with PDND patients without orthostatic diastolic hypotension ($p = 0.029$ and $p = 0.014$) or with controls ($p = 0.004$ and $p = 0.005$). The ratios did not differ between PDND patients without orthostatic diastolic hypotension and controls.

Cohort 1: Angiogenesis biomarkers and CSF/plasma

albumin ratio. Increased levels of CSF/plasma albumin ratio often reflects increased permeability of the BBB.¹⁸ The CSF/plasma albumin ratio differed between the diagnostic groups ($p = 0.005$; table 1). The ratio was higher in PDND patients ($p = 0.001$) and PDD patients ($p = 0.027$) compared to controls (figure 2A), while no difference was found between PDND patients and PDD patients. In all patients with PD, higher CSF/plasma albumin ratio was associated with higher CSF levels of VEGF ($\beta = 0.48$, $p < 0.001$; figure 2B) and PIGF

Figure 2 Angiogenesis biomarkers, blood-brain barrier, and WMLs in controls, PDND, and PDD (cohort 1)

(A) The CSF/plasma albumin ratio is higher in patients with Parkinson disease without dementia (PDND) and patients with Parkinson disease with dementia (PDD) than in controls. (B–E) Vascular endothelial growth factor (VEGF) and placental growth factor (PIGF) are associated with the CSF/plasma albumin ratio and white matter lesions (WMLs) in patients with Parkinson disease. Angiogenesis biomarkers were skewed and therefore ln-transformed before statistical analyses; p values are from analysis of covariance (A) and from linear regression (B–E) adjusting for age and sex as described in the Methods; data in (A) are presented mean $\pm$ 95% confidence interval.

($\beta = 0.43$, $p < 0.001$; figure 2C), sVEGFR-2 ($\beta = 0.41$, $p < 0.001$), and IL-8 ($\beta = 0.27$, $p = 0.009$), as well as with higher VEGF/sVEGFR1 ratio ($\beta = 0.33$, $p = 0.001$) and PlGF/sVEGFR1 ratio ($\beta = 0.29$, $p = 0.006$). We observed similar associations when only including PDND patients (data not shown). In the control group, we did not find any significant associations between CSF/plasma albumin ratio and any of the angiogenesis variables or ratios (all $p > 0.16$).

Cohort 1: Angiogenesis biomarkers, WMLs, and CMB. WMLs (total age-related white matter changes [ARWMC] score) did not differ between the diagnostic groups (table 1). In patients with PD, the total burden of WMLs was associated with higher CSF levels of VEGF ($\beta = 0.40$, $p = 0.004$; figure 2D), PlGF ($\beta = 0.42$, $p = 0.002$; figure 2E), and IL-8 ($\beta = 0.24$, $p = 0.044$), as well as with higher VEGF/sVEGFR-1 ratio ($\beta = 0.37$, $p = 0.004$) and PlGF/sVEGFR-1 ratio ($\beta = 0.38$, $p = 0.003$). When only

including PDND patients, we found similar results (data not shown). In the control group, we did not find any significant associations between total ARWMC score and any of the angiogenesis variables or ratios (all $p > 0.14$).

Patients with PD with signs of CMB ($n = 13$) had higher VEGF ($p = 0.051$) and PlGF ($p = 0.020$) levels than those who did not have CMB ($n = 37$).

Cohort 1: Angiogenesis biomarkers and markers of neurodegeneration and neuroinflammation. NFL is a marker of axonal neurodegeneration.¹⁹ In patients with PD, higher levels of NFL in CSF are associated with more severe disease.²⁰ Positive associations were found between CSF NFL levels and VEGF ($\beta = 0.47$, $p = 0.001$) and PlGF ($\beta = 0.31$, $p = 0.021$), as well as CSF/plasma albumin ratio ($\beta = 0.41$, $p = 0.016$) in patients with PD but not in controls. Supporting the role of NFL in axonal degeneration, we observed a strong association between WMLs (total ARWMC score) and CSF NFL levels ($\beta = 0.60$, $p = 0.003$).

CSF t-tau levels are considered to reflect cortical neurodegeneration,¹⁸ and the CSF tau levels did not correlate with PlGF or VEGF. Further, there were no correlations between PlGF and VEGF and the Alzheimer-specific biomarkers CSF p-tau or A β 42.

To study possible associations of angiogenesis biomarkers with neuroinflammation, we measured the CSF levels of MCP-1, which is a chemoattractant expressed by microglia and astrocytes in the brain²¹ that can be reliably measured in CSF.²² MCP-1 was associated with VEGF in the patients with PD ($\beta = 0.25$, $p = 0.027$) but not in the controls.

The associations with markers of neurodegeneration and neuroinflammation were similar when only including PDND patients (data not shown).

Cohort 2: Validation study. In this cohort of controls and PDD/DLB patients, we analyzed CSF levels of VEGF, PlGF, and sVEGFR1. The patient group displayed higher mean levels of PlGF ($p = 0.002$), VEGF/sVEGFR-1 ratio ($p < 0.001$), and PlGF/sVEGFR-1 ratio ($p < 0.001$) compared to the healthy controls (figure 3, A–C). In this cohort of

patients we also confirmed the association of CSF/plasma albumin ratio with VEGF ($\beta = 0.48$, $p < 0.001$) and PlGF ($\beta = 0.42$, $p = 0.002$). Using VEGF V-PLEX kit, we found that VEGF levels were elevated in the PDD/DLB patients compared to the healthy controls (supplemental data).

Cohort 3: Validation study. In this cohort of neuropathologically confirmed cases, we measured VEGF, PlGF, and sVEGFR-1 in postmortem ventricular CSF samples from PDND and PDD participants as well as healthy controls. In agreement with our previous findings, both the PDND and PDD groups showed elevated levels of VEGF ($p = 0.016$ and $p = 0.034$) and PlGF ($p = 0.037$ and $p = 0.002$) compared to the controls (figure 3, D and E). We did not find any difference in these analytes between the PDND and PDD groups. In all patients with PD, the total burden of WMLs was associated with higher CSF levels of VEGF ($\beta = 0.24$, $p = 0.016$), confirming our previous finding when comparing WMLs according to MRI and angiogenesis biomarkers in CSF. When only including the

Figure 3 CSF levels of angiogenesis biomarkers (cohorts 2 and 3)

(A–C) Placental growth factor (PlGF), vascular endothelial growth factor (VEGF)/sVEGFR-1 ratio, and PlGF/sVEGFR-1 ratio are higher in patients with Parkinson disease with dementia (PDD)/patients with dementia with Lewy bodies (DLB) than in controls in cohort 2. (D, E) VEGF and PlGF levels are elevated in patients with Parkinson disease without dementia (PDND) and patients with PDD compared to controls in cohort 3 with neuropathologically confirmed diagnosis. Angiogenesis biomarkers were skewed and therefore ln-transformed before statistical analyses; p values are from analysis of covariance adjusting for age and sex as described in the Methods; data are presented as mean $\pm$ 95% confidence interval.

PDND patients, we found similar results (data not shown).

There were no associations between CSF levels of either VEGF or PlGF and amyloid plaque density (quantified according to the Consortium to Establish a Registry for Alzheimer's Disease templates) or Braak tangle scores in patients with PD, suggesting that changes in angiogenic factors are not related to AD amyloid or tau pathology. These findings confirm the results obtained in cohort 1 using CSF AD biomarkers.

DISCUSSION Using a well-characterized clinical sample, this study demonstrates that CSF levels of VEGF, PlGF, and sVEGFR-2 are high, while levels of Ang2 are low, in PDND and PDD patients compared to non-PD controls. We corroborate our findings of elevated CSF levels of VEGF and PlGF in 2 additional cohorts of patients with PD with and without dementia.

The increased CSF levels of angiogenesis factors in PD are in keeping with postmortem studies demonstrating high levels of VEGF and elevated numbers of endothelial cell nuclei and vessels in the SNpc of patients with PD, as well as primates with parkinsonian lesions.^{1–3,23} Previous reports have also suggested that PD might be accompanied by BBB dysfunction.^{24,25} In agreement with these reports, we found higher CSF/plasma albumin ratio in patients with PD compared to controls. Furthermore, we observed robust positive correlations between angiogenic factors and CSF/plasma albumin ratio, which is interesting considering that VEGF is a potent trigger of vascular leakage and BBB dysfunction.^{26,27} In other neurodegenerative diseases, mild but chronic increases in BBB permeability are known to contribute to neuronal dysfunction and neurodegeneration.²⁸ Of relevance to PD, intranigral administration of VEGF induces BBB breakdown and loss of dopaminergic neurons.²⁹ On the contrary, the angiogenic factor Ang2 decreases BBB leakage.³⁰ Thus, increased levels of VEGF and PlGF, and decreased levels of Ang2, may concur to produce BBB dysfunction and neuronal toxicity in the affected brain regions. Accordingly, we found strong correlations between the CSF levels of NFL (a marker of axonal degeneration) and angiogenesis biomarkers in patients with PD.

BBB abnormalities are likely to be associated with structurally altered and leaky small vessels that may result in CMB.³¹ High serum VEGF and increased vascular permeability have been previously linked to CMB in acute ischemic stroke.³² Here we show that patients with PD with CMB have elevated CSF levels of VEGF and PlGF. These findings raise the possibility that angiogenic factors are involved in

the pathogenesis of CMB, in addition to BBB dysfunction, in PD.

The mechanisms behind the observed alterations in angiogenesis biomarkers in PD are yet to be elucidated. In the present study, we found that patients with PD with orthostatic diastolic hypotension have increased levels of VEGF and PlGF compared to patients without orthostatic diastolic hypotension. Orthostatic hypotension might result in recurrent episodes of transient cerebral hypoperfusion and hypoxia in cases with suboptimal autoregulation of cerebral blood flow as for example in PD.^{33,34} Hypoxia-induced VEGF signaling is one of the main molecular pathways triggering angiogenic responses.³⁵ Hypoperfusion and hypoxia have been also proposed as a potential mechanism underlying previously observed association between orthostatic hypotension and WMLs.^{16,36} We found that increased levels of VEGF and PlGF in the CSF of patients with PD were related to more pronounced WMLs. Thus, elevated levels of angiogenesis biomarkers in patients with PD in our study may partly depend on the hypoxic environment associated with orthostatic hypotension and WMLs. WMLs have been linked to gait abnormalities and postural instability in large cohorts of healthy elderly individuals as well as patients with PD.³⁷ Accordingly, in our PD cohorts elevated levels of VEGF and PlGF correlated with poor performance on the Tandem Gait Test. These findings suggest that WMLs and increased angiogenesis underpin, at least to some extent, the treatment-resistant symptoms that develop during the course of PD, such as postural instability and gait difficulties.

In our study, VEGF positively correlated with MCP-1, a proinflammatory chemokine expressed by astrocytes and microglia.²¹ Interestingly, one previous study reported a greater number of activated microglia together with signs of angiogenesis in the SNpc of parkinsonian brain.⁴ Others have shown that VEGF production can be triggered in astrocytes by proinflammatory cytokines.^{38,39} Collectively, these data suggest a possible link between neuroinflammation and angiogenesis in PD.

One potential limitation of the current study is the cross-sectional study design, thus causality may not be inferred. Another limitation is that it is not possible to obtain any regional or structural information using CSF. On the other hand, analysis of material, such as CSF, from living individuals with PD has some advantages over postmortem investigations. Relatively large series of patients can be studied, findings can easily be related to current symptomatology and imaging data in the same participants, and the results are not biased by death-related events, such as hypoxia. The proportion of

individuals with major somatic comorbidities or using of anti-inflammatory medications did not significantly differ across groups, thus these factors are unlikely to have confounded our findings. A strength of the present study was the multimodal characterization of the original prospective cohort with clinical assessments, brain imaging, and different CSF biomarkers. Further, the main results obtained in the original cohort were validated in 2 independent cohorts, of which one consisted of neuropathologically confirmed cases. Finally, different ELISAs were used to measure VEGF in CSF, resulting in similar differences between diagnostic groups.

In the present study, we established that angiogenic factors are upregulated in the CSF in PD. The angiogenic activation was associated with gait difficulties, BBB dysfunction, WMLs, CBM, neuroinflammation, and axonal degeneration in patients but not in the control group, suggesting that these interactions are specific for PD. Our results warrant further mechanistic studies on the role of angiogenesis and cerebrovascular dysfunction in the development of some symptoms that respond poorly to dopamine replacement therapy in PD. These studies may have an important impact on the design of new personalized therapeutic approaches.

AUTHOR CONTRIBUTIONS

S.J. codesigned the study; collected, analyzed, and interpreted the data; conducted literature searches; prepared figures; and cowrote the manuscript. D.L. codesigned the study, analyzed and interpreted the data, conducted literature searches, and cowrote the manuscript. V.F., S.H., H.Z., K.B., C.H.A., T.G.B., G.S., D.v.W., and E.L. collected and analyzed the data and reviewed the manuscript for intellectual content. M.A.C. codesigned the study, interpreted the data, conducted literature searches, overviewed data collection, cowrote the manuscript, and obtained funding. O.H. was the principal designer and coordinator of the study; overviewed collection, analysis, and interpretation of the study data; cowrote the manuscript; and obtained funding.

ACKNOWLEDGMENT

The authors thank the research nurses, Ann Johansson, Jan Reimer, and Katarina Johansson, for their assistance in the clinical collection of CSF samples and data from study participants and healthy controls; and The Arizona Parkinson's Disease Consortium and the Banner Sun Health Research Institute Brain and Body Donation Program of Sun City, Arizona, for providing human CSF.

STUDY FUNDING

The Brain and Body Donation Program is supported by the National Institute of Neurological Disorders and Stroke (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), the Arizona Department of Health Services (contract 211002, Arizona Alzheimer's Research Center), the Arizona Biomedical Research Commission (contracts 4001, 0011, 05-901 and 1001 to the Arizona Parkinson's Disease Consortium), and the Michael J. Fox Foundation for Parkinson's Research. The study was supported by the European Research Council, the Swedish Research Council, the Parkinson Foundation of Sweden, the Michael J. Fox Foundation, the Crafoord Foundation, the Swedish Brain Foundation, the Swedish Federal Government under the ALF Agreement, The Torsten Söderberg Foundation, Multipark, and the

Knut and Alice Wallenberg Foundation. The funding sources had no role in the design and conduct of the study; in the collection, analysis, or interpretation of the data; or in the preparation, review, or approval of the manuscript.

DISCLOSURE

S. Janelidze, D. Lindqvist, V. Francardo, S. Hall, and H. Zetterberg report no disclosures relevant to the manuscript. K. Blennow has served on advisory boards for IBL International, Pfizer, Roche Diagnostics, Lilly, and Kyowa Kirin Pharma. C. Adler, T. Beach, G. Serrano, D. van Westen, E. Londos, M. Cenci, and O. Hansson report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Received March 25, 2015. Accepted in final form July 9, 2015.

REFERENCES

1. Faucheux BA, Bonnet AM, Agid Y, Hirsch EC. Blood vessels change in the mesencephalon of patients with Parkinson's disease. *Lancet* 1999;353:981-982.
2. Wada K, Arai H, Takanashi M, et al. Expression levels of vascular endothelial growth factor and its receptors in Parkinson's disease. *Neuroreport* 2006;17:705-709.
3. Barcia C, Bautista V, Sanchez-Bahillo A, et al. Changes in vascularization in substantia nigra pars compacta of monkeys rendered parkinsonian. *J Neural Transm* 2005;112:1237-1248.
4. Desai Bradaric B, Patel A, Schneider JA, Carvey PM, Hendey B. Evidence for angiogenesis in Parkinson's disease, incidental Lewy body disease, and progressive supranuclear palsy. *J Neural Transm* 2012;119:59-71.
5. Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med* 2003;9:669-676.
6. Fiedler U, Augustin HG. Angiopoietins: a link between angiogenesis and inflammation. *Trends Immunol* 2006;27:552-558.
7. Kiefer F, Siekmann AF. The role of chemokines and their receptors in angiogenesis. *Cell Mol Life Sci* 2011;68:2811-2830.
8. Gelb DJ, Oliver E, Gilman S. Diagnostic criteria for Parkinson disease. *Arch Neurol* 1999;56:33-39.
9. Emre M, Aarsland D, Brown R, et al. Clinical diagnostic criteria for dementia associated with Parkinson's disease. *Mov Disord* 2007;22:1689-1707; quiz 1837.
10. Hertz J, Nagga K, Minthon L, Hansson O. Changes in cerebrospinal fluid and blood plasma levels of IGF-II and its binding proteins in Alzheimer's disease: an observational study. *BMC Neurol* 2014;14:64.
11. Lindqvist D, Hall S, Surova Y, et al. Cerebrospinal fluid inflammatory markers in Parkinson's disease: associations with depression, fatigue, and cognitive impairment. *Brain Behav Immun* 2013;33:183-189.
12. Zetterberg H, Jakobsson J, Redsater M, et al. Blood-cerebrospinal fluid barrier dysfunction in patients with bipolar disorder in relation to antipsychotic treatment. *Psychiatry Res* 2014;217:143-146.
13. Ambati BK, Nozaki M, Singh N, et al. Corneal avascularity is due to soluble VEGF receptor-1. *Nature* 2006;443:993-997.
14. Kendall RL, Thomas KA. Inhibition of vascular endothelial cell growth factor activity by an endogenously encoded soluble receptor. *Proc Natl Acad Sci USA* 1993;90:10705-10709.
15. Qi JW, Qin TT, Xu LX, et al. TNFSF15 inhibits vasculogenesis by regulating relative levels of membrane-bound and soluble isoforms of VEGF receptor 1. *Proc Natl Acad Sci USA* 2013;110:13863-13868.

16. Kruit MC, Thijs RD, Ferrari MD, Launer LJ, van Buchem MA, van Dijk JG. Syncope and orthostatic intolerance increase risk of brain lesions in migraineurs and controls. *Neurology* 2013;80:1958–1965.
17. Roman GC, Erkinjuntti T, Wallin A, Pantoni L, Chui HC. Subcortical ischaemic vascular dementia. *Lancet Neurol* 2002;1:426–436.
18. Blennow K, Hampel H, Weiner M, Zetterberg H. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol* 2010;6:131–144.
19. Skillback T, Zetterberg H, Blennow K, Mattsson N. Cerebrospinal fluid biomarkers for Alzheimer disease and subcortical axonal damage in 5,542 clinical samples. *Alzheimers Res Ther* 2013;5:47.
20. Hall S, Ohrfelt A, Constantinescu R, et al. Accuracy of a panel of 5 cerebrospinal fluid biomarkers in the differential diagnosis of patients with dementia and/or parkinsonian disorders. *Arch Neurol* 2012;69:1445–1452.
21. Yao Y, Tsirka SE. Monocyte chemoattractant protein-1 and the blood-brain barrier. *Cell Mol Life Sci* 2014;71:683–697.
22. Westin K, Buchhave P, Nielsen H, Minthon L, Janciauskiene S, Hansson O. CCL2 is associated with a faster rate of cognitive decline during early stages of Alzheimer's disease. *PLoS One* 2012;7:e30525.
23. Ohlin KE, Francardo V, Lindgren HS, et al. Vascular endothelial growth factor is upregulated by L-dopa in the parkinsonian brain: implications for the development of dyskinesia. *Brain* 2011;134:2339–2357.
24. Kortekaas R, Leenders KL, van Oostrom JC, et al. Blood-brain barrier dysfunction in parkinsonian midbrain in vivo. *Ann Neurol* 2005;57:176–179.
25. Pisani V, Stefani A, Pierantozzi M, et al. Increased blood-cerebrospinal fluid transfer of albumin in advanced Parkinson's disease. *J Neuroinflammation* 2012;9:188.
26. Argaw AT, Asp L, Zhang J, et al. Astrocyte-derived VEGF-A drives blood-brain barrier disruption in CNS inflammatory disease. *J Clin Invest* 2012;122:2454–2468.
27. Weis SM, Cheres DA. Pathophysiological consequences of VEGF-induced vascular permeability. *Nature* 2005;437:497–504.
28. Zlokovic BV. The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron* 2008;57:178–201.
29. Rite I, Machado A, Cano J, Venero JL. Blood-brain barrier disruption induces in vivo degeneration of nigral dopaminergic neurons. *J Neurochem* 2007;101:1567–1582.
30. Marteau L, Valable S, Divoux D, et al. Angiotensin-2 is vasoprotective in the acute phase of cerebral ischemia. *J Cereb Blood Flow Metab* 2013;33:389–395.
31. Yates PA, Villemagne VL, Ellis KA, Desmond PM, Masters CL, Rowe CC. Cerebral microbleeds: a review of clinical, genetic, and neuroimaging associations. *Front Neurol* 2014;4:205.
32. Dassan P, Brown MM, Gregoire SM, Keir G, Werring DJ. Association of cerebral microbleeds in acute ischemic stroke with high serum levels of vascular endothelial growth factor. *Arch Neurol* 2012;69:1186–1189.
33. Novak V, Novak P, Spies JM, Low PA. Autoregulation of cerebral blood flow in orthostatic hypotension. *Stroke* 1998;29:104–111.
34. Vokatch N, Grotzsch H, Mermillod B, Burkhard PR, Sztajzel R. Is cerebral autoregulation impaired in Parkinson's disease? A transcranial Doppler study. *J Neurol Sci* 2007;254:49–53.
35. Kim YW, Byzova TV. Oxidative stress in angiogenesis and vascular disease. *Blood* 2014;123:625–631.
36. Oh YS, Kim JS, Lee KS. Orthostatic and supine blood pressures are associated with white matter hyperintensities in Parkinson disease. *J Mov Disord* 2013;6:23–27.
37. Bohnen NI, Albin RL. White matter lesions in Parkinson disease. *Nat Rev Neurol* 2011;7:229–236.
38. Argaw AT, Zhang Y, Snyder BJ, et al. IL-1 β regulates blood-brain barrier permeability via reactivation of the hypoxia-angiogenesis program. *J Immunol* 2006;177:5574–5584.
39. Loeffler S, Fayard B, Weis J, Weissenberger J. Interleukin-6 induces transcriptional activation of vascular endothelial growth factor (VEGF) in astrocytes in vivo and regulates VEGF promoter activity in glioblastoma cells via direct interaction between STAT3 and Sp1. *Int J Cancer* 2005;115:202–213.

Experience the Excellence of the 2016 AAN Annual Meeting

Registration is now open for the 68th AAN Annual Meeting, set to take place Friday, April 15, through Thursday, April 21, 2016, at the Vancouver Convention Centre in Vancouver, BC, Canada. Experience the excellence of new, exciting changes, including: one low, single registration rate; more than 230 education programs in two-hour increments; plenary sessions every day starting Friday; scientific poster sessions every day starting Saturday; interactive and experiential learning and networking opportunities all week long, and more! The money-saving early registration deadline is March 24, 2016. Visit AAN.com/view/AM16 and register today!

Neurology[®]

Increased CSF biomarkers of angiogenesis in Parkinson disease

Shorena Janelidze, Daniel Lindqvist, Veronica Francardo, et al.

Neurology 2015;85:1834-1842 Published Online before print October 28, 2015

DOI 10.1212/WNL.0000000000002151

This information is current as of October 28, 2015

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2015 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/85/21/1834.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2015/10/28/WNL.0000000000002151.DC1 http://n.neurology.org/content/suppl/2015/10/28/WNL.0000000000002151.DC2
References	This article cites 39 articles, 6 of which you can access for free at: http://n.neurology.org/content/85/21/1834.full#ref-list-1
Citations	This article has been cited by 6 HighWire-hosted articles: http://n.neurology.org/content/85/21/1834.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Cerebrospinal Fluid http://n.neurology.org/cgi/collection/cerebrospinal_fluid MRI http://n.neurology.org/cgi/collection/mri Parkinson's disease with dementia http://n.neurology.org/cgi/collection/parkinsons_disease_with_dementia Parkinson's disease/Parkinsonism http://n.neurology.org/cgi/collection/parkinsons_disease_parkinsonism
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2015 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.