Teaching NeuroImages:

Noninfectious cyst as an unusual complication of deep brain stimulation

Harsh V. Gupta, MD
Mark K. Lyons, MD
Shyamal H. Mehta, MD,
PhD

Correspondence to
Dr. Gupta:
dr.harshgupta@gmail.com

Figure 1 Axial fluid-attenuated inversion recovery (FLAIR), T2, postcontrast T1, and diffusion-weighted imaging (DWI)

(A) Axial FLAIR: Cyst is seen adjacent to the DBS lead with surrounding edema. (B) Axial T2: Cyst is visualized adjacent to the DBS lead with vasogenic edema. (C) T1 postcontrast: Absence of contrast enhancement. (D) Axial DWI: Absence of restricted diffusion.

A 73-year-old woman with essential tremor complained of cognitive difficulties, right-sided weakness, and balance problems 3 months after bilateral deep brain stimulation (DBS) surgery. MRI brain showed a cyst adjacent to the DBS electrode on the left side along with vasogenic edema (figure 1). CSF and blood studies did not demonstrate any evidence of infection. Dexamethasone initiation led to a gradual improvement in symptoms and resolution of edema (figure 2).

The tremor did not disappear after cyst development and a higher voltage was required on the left side to control the tremor. Removal of the cyst should be considered when it continues to grow or leads to

undesirable symptoms or lack of therapeutic effect from the DBS.^{1,2}

AUTHOR CONTRIBUTIONS

Dr. Gupta and Dr. Mehta: drafting of the manuscript, examination of the patient, and review of the imaging findings. Dr. Lyons: examination of the patient and review of the images.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

H. Gupta and M. Lyons report no disclosures relevant to the manuscript. S. Mehta has consulted with Allergan Inc., US World Meds, Merz Pharma, and Cynapsus Therapeutics. Go to Neurology.org for full disclosures.

Download teaching slides:
Neurology.org

From the Departments of Neurology (H.V.G., S.H.M.) and Neurosurgery (M.K.L.), Mayo Clinic, Scottsdale, AZ.

Figure 2 Axial T2 and fluid-attenuated inversion recovery (FLAIR)

(A) Axial T2: Resolution of the surrounding edema and minimal decrease in cyst size. (B) Axial FLAIR: Resolution of the surrounding edema and minimal decrease in cyst size.

REFERENCES

1. Ramirez-Zamora A, Levine D, Sommer DB, Dalfino J, Novak P, Pilitsis JG. Intraparenchymal cyst development after deep brain stimulator placement. *Stereotact Funct Neurosurg* 2013;91:338–341.
2. Sharma VD, Bona AR, Mantovani A, et al. Cystic lesions as a rare complication of deep brain stimulation. *Mov Disord Clin Pract* 2015;3:87–90.

Neurology®

Teaching *NeuroImages*: Noninfectious cyst as an unusual complication of deep brain stimulation

Harsh V. Gupta, Mark K. Lyons and Shyamal H. Mehta

Neurology 2016;87:e223-e224

DOI 10.1212/WNL.0000000000003284

This information is current as of October 31, 2016

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/87/18/e223.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2016/10/31/WNL.0000000000003284.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/87/18/e223.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): MRI http://n.neurology.org/cgi/collection/mri Surgery/Stimulation http://n.neurology.org/cgi/collection/surgery-stimulation
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2016 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.