5. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;55:181–184.
6. Postuma RB, Berg D, Stern M, et al. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord* 2015;30:1591–1601.
7. Parnetti L, Chiasserini D, Persichetti E, et al. Cerebrospinal fluid lysosomal enzymes and alpha-synuclein in Parkinson's disease. *Mov Disord* 2014;29:1019–1027.
8. Adler CH, Dugger BN, Hentz JG, et al. Peripheral synucleinopathy in early Parkinson's disease: submandibular gland needle biopsy findings. *Mov Disord* 2016;31:250–256.
9. Donadio V, Incensi A, Piccinini C, et al. Skin nerve misfolded α -synuclein in pure autonomic failure and Parkinson disease. *Ann Neurol* 2016;79:306–316.

PRACTICE GUIDELINE UPDATE SUMMARY: CORTICOSTEROID TREATMENT OF DUCHENNE MUSCULAR DYSTROPHY: REPORT OF THE GUIDELINE DEVELOPMENT SUBCOMMITTEE OF THE AMERICAN ACADEMY OF NEUROLOGY

Haluk Topaloglu, Ankara, Turkey: I read with interest the guidelines on corticosteroid treatment in Duchenne muscular dystrophy (DMD).¹ As a pediatrician, I find 0.75 mg/kg/d of prednisone very high. I understand the principle of initiating with a higher dose; however, maintaining this regimen beyond weeks is not possible in practice. The authors also quoted a time frame of 6 months. We do not even give this amount (cumulatively) of steroids to patients with juvenile dermatomyositis. Of course, all is a matter of tailoring the drug, but this should be done early. Serious side effects are virtually inevitable at this recommended dose, which is cumulative and naturally will create other health issues (sometimes irreversible). New therapies (molecular) are approaching and, in the meantime, we face the dilemma of keeping the patient in good health while fearing that once

walking is lost it will not come back. I realize this fact. However, the general health of the patient would be a concern.

Author Response: David Gloss, Charleston, WV; Richard T. Moxley III, Rochester, NY; Stephen Ashwal, Loma Linda, CA; Maryam Oskoui, Montreal, Canada: We thank Dr. Topaloglu for sharing his personal experience. The side effects of long-term steroids are substantial and should be part of the decision-making process.

The evidence supports 0.75 mg/kg/d as the preferred dose of prednisone in this patient population.¹ There is robust short- and long-term evidence of prednisone's efficacy in motor function, pulmonary function, and cardiac function. In practice, patients with DMD have been maintained on this dose of prednisone for years. Daily corticosteroid use is considered by many in the field to be standard of care. Furthermore, there is evidence that a lower daily dose at 0.3 mg/kg/d, although associated with fewer side effects, is less effective.²

New therapies are on the horizon. However, they are not currently clinically available. Most clinical trials on these new therapies also consider daily steroid therapy as standard of care, making it a criterion for study entry in many cases. In our next update, if these new therapies are available, we plan to include them.

© 2016 American Academy of Neurology

1. Gloss D, Moxley RT III, Ashwal S, Oskoui M. Practice guideline update summary: Corticosteroid treatment of Duchenne muscular dystrophy: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology* 2016;86:465–472.
2. Griggs RC, Moxley RT III, Mendell JR, et al. Prednisone in Duchenne dystrophy: a randomized, controlled trial defining the time course and dose response: clinical investigation of Duchenne dystrophy group. *Arch Neurol* 1991;48:383–388.

CORRECTION

NDUFA-1 is not a nuclear modifier gene in Leber hereditary optic neuropathy

In the Clinical/Scientific Note “NDUFA-1 is not a nuclear modifier gene in Leber hereditary optic neuropathy” by P. Yu-Wai-Man et al.,¹ there is an error in the first author's name, which should have read “P. Yu-Wai-Man” rather than “P.Y.W. Man” as originally published.

REFERENCE

1. Yu-Wai-Man P, Brown DT, Wehnert MS, et al. NDUFA-1 is not a nuclear modifier gene in Leber hereditary optic neuropathy. *Neurology* 2002;58:1861–1862.

Author disclosures are available upon request (journal@neurology.org).

Neurology[®]

NDUFA-1 is not a nuclear modifier gene in Leber hereditary optic neuropathy

Neurology 2016;87;238

DOI 10.1212/WNL.0000000000002857

This information is current as of July 11, 2016

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/87/2/238.2.full>

References

This article cites 1 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/87/2/238.2.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2016 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.