Mystery Case: Brown-Séquard syndrome caused by idiopathic spinal cord herniation

Erich Talamoni Fonoff,
MD, PhD
William Omar Contreras
Lopez, MD, PhD
Manoel Jacobsen Teixeira,
MD, PhD

Correspondence to
Dr. Fonoff:
fonoffet@usp.br

Figure T2-weighted MRI of the thoracic spine shows signs of spinal cord herniation

T2-weighted MRI of the thoracic spine displays anterolateral spinal cord deformation in the coronal (A), axial (B), and sagittal planes (C); this image corresponds to a spinal cord kink between the 6th and 7th thoracic vertebrae and a secondary expansion of the subarachnoid space. Axial image shows part of the cord protruded beyond the dural limits in anterolateral quadrant of spinal canal.¹

A 48-year-old man developed numbness in the left leg, which progressed gradually to paresis and urinary incontinence. Neurologic examination revealed a left Brown-Séquard syndrome with leg paresis, mild spasticity, reduced proprioception, and contralateral thermal and painful hypoesthesia below T6. MRI revealed a thoracic spinal cord herniation (SCH) (figure). Idiopathic SCH is relatively rare. Pathogenesis involves a dura mater defect (see video on the *Neurology*[®] Web site at Neurology.org); herniation develops over a progressive pressure gradient through the dural fissure.² Surgical reduction is typically performed if symptoms progress, but mild symptoms may be eligible for conservative treatment and monitoring. Surgical spinal reduction and dural repair usually reverses neurologic deficits.

AUTHOR CONTRIBUTIONS

Erich Talamoni Fonoff: drafting/revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and final approval, acquisition of data. William Omar Contreras Lopez: drafting/revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and final approval, acquisition of data. Manoel Jacobsen Teixeira: drafting/revising the manuscript, accepts responsibility for conduct of research and final approval, study supervision.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

REFERENCES

1. Corbetta M, Capone F, Denaro V, et al. Clinical reasoning: a case of slowly progressive painful paraparesis. *Neurology* 2014;82:1476–1479.
2. Najjar MW, Baesa SS, Lingawi SS. Idiopathic spinal cord herniation: a new theory of pathogenesis. *Surg Neurol* 2004; 62:161–170; discussion 170–171.

MYSTERY CASE RESPONSES

The Mystery Case series was initiated by the *Neurology*[®] Resident & Fellow Section to develop the clinical reasoning skills of trainees. Residency programs, medical student preceptors, and individuals were invited to use this Mystery Case as an educational tool. Responses were solicited through a group e-mail sent to the American Academy of Neurology Consortium of Neurology Residents and Fellows and through social media.

All the responses came from individuals. Thirty-three percent correctly identified the patient's presentation as being a Brown-Séquard syndrome. Sixty-seven percent of the respondents correctly interpreted the MRI findings as spinal cord herniation. Thirty-three percent suggested a diagnosis of dural

Supplemental data
at Neurology.org

From the Division of Functional Neurosurgery, Department of Neurology, University of São Paulo Medical School, Brazil.

meningioma—while this is an important item on the differential diagnosis, the MRI reveals deformation of normal cord tissue rather than abnormal tissue growth as one would see with a meningioma. This can be seen especially well in the sagittal section.

The most complete response was provided by Dr. Jeremy Cutsforth-Gregory from the Mayo Clinic, who provided a clear localization for the patient's symptoms—specifically the involvement of the left

lateral corticospinal tract and left dorsal column, and the right spinothalamic tract—and recognized the MRI findings as representing an idiopathic spinal cord herniation.

Aravind Ganesh, MD

Department of Clinical Neurosciences, University of Calgary, Canada; and Centre for Prevention of Stroke & Dementia, Nuffield Department of Clinical Neurosciences, University of Oxford, UK

Neurology®

Mystery Case: Brown-Séquard syndrome caused by idiopathic spinal cord herniation

Erich Talamoni Fonoff, William Omar Contreras Lopez and Manoel Jacobsen Teixeira

Neurology 2016;87:e34-e35

DOI 10.1212/WNL.0000000000002886

This information is current as of July 25, 2016

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/87/4/e34.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2016/07/24/WNL.0000000000002886.DC1
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/87/4/e34.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord Clinical neurology examination http://n.neurology.org/cgi/collection/clinical_neurology_examination MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2016 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.