Teaching *NeuroImages*: Homotopic motor distribution on fMRI in closed-lip schizencephaly

Luca Bartolini, MD
Amy Kao, MD
Ashley Zachery, BA
Jonathan G. Murnick,
MD
William D. Gaillard, MD

Correspondence to
Dr. Bartolini:
LBartoli@childrensnational.org

Figure 1 3T brain MRI and functional MRI of a 16-year-old boy with intractable focal epilepsy due to schizencephaly and polymicrogyria

Axial T1-weighted (left) and axial and coronal T2-weighted (right) images demonstrate polymicrogyria extending from the cerebral convexity to the ventricular margin. Focal cleft where abnormal gray matter meets the ependymal surface is consistent with closed-lip schizencephaly (A). Activation with repetitive bilateral foot (B), bilateral finger (C), and tongue tapping (D) reveals homotopic distribution of motor cortex with hand deep into the cleft centrally, foot mesially, and mouth laterally at the edge of the area of schizencephaly. The threshold for the study was set at top 5% of voxels.

Download teaching slides:
Neurology.org

From the Center for Neuroscience and Behavioral Medicine (L.B., A.K., A.Z., W.D.G.) and the Division of Diagnostic Imaging and Radiology (J.G.M.), Children's National Health System, The George Washington University, Washington, DC.

Go to Neurology.org for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Figure 2 Skull radiograph shows depth electrode placement

Motor mapping is shown as red dots next to the electrodes. Right lateral (RL): stimulation of electrodes 1-2 resulted in motion of the left hand; right medial (RM): stimulation of electrodes 1-2 resulted in left arm and face motion; 3-4, jaw motion; 5-6, left side of mouth motion; 7-8, left knee twitch; left lateral (LL): stimulation of electrodes 1-2 resulted in right hand then right arm then right leg twitch; 3-4, right leg extension; 5, right foot twitch.

A 16-year-old boy with intractable focal epilepsy with preserved awareness and attention-deficit/hyperactivity disorder underwent evaluation for epilepsy surgery. Epilepsy onset occurred at age 5 years; seizures arise from sleep and are characterized by right arm extension with preserved consciousness. Ictal EEG has no clear localization or lateralization. MRI brain demonstrates

bifrontal closed-lip schizencephaly. Functional MRI (fMRI) shows homotopic distribution of tongue/finger/foot motor function at the edge of and along the depth of the schizencephalic cleft, supported by intracranial mapping (figures 1 and 2). fMRI is a reliable tool to identify the eloquent cortex in the planning for epilepsy surgery.^{1,2}

AUTHOR CONTRIBUTIONS

Dr. Bartolini designed and drafted the article. Dr. Kao contributed to the design and draft of the article. A. Zachery processed the fMRI and contributed to the article draft. Dr. Murnick reviewed the manuscript and contributed to the interpretation of the structural MRI. Dr. Gaillard reviewed the manuscript and contributed to the interpretation of the fMRI.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

L. Bartolini serves as the Practice Current Section Editor of *Neurology*[®] *Clinical Practice* and is a member of the editorial team of the *Neurology Resident & Fellow* section. A. Kao, A. Zachery, and J. Murnick report no disclosures relevant to the manuscript. W. Gaillard is Associate Editor of *Epilepsia* and *Epilepsy Research*. Go to Neurology.org for full disclosures.

REFERENCES

1. Martinez-Rios C, McAndrews MP, Logan W, Krings T, Lee D, Widjaja E. MRI in the evaluation of localization-related epilepsy. *J Magn Reson Imaging* 2016;44:12-22.
2. Gaillard WD, Berl MM. Functional magnetic resonance imaging: functional mapping. *Handb Clin Neurol* 2012; 107:387-398.

Neurology®

Teaching *NeuroImages*: Homotopic motor distribution on fMRI in closed-lip schizencephaly

Luca Bartolini, Amy Kao, Ashley Zachery, et al.

Neurology 2017;88:e24-e25

DOI 10.1212/WNL.0000000000003493

This information is current as of January 9, 2017

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/88/2/e24.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2017/01/09/WNL.0000000000003493.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/88/2/e24.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Pediatric http://n.neurology.org/cgi/collection/all_pediatric Cortical localization http://n.neurology.org/cgi/collection/cortical_localization Epilepsy surgery http://n.neurology.org/cgi/collection/epilepsy_surgery Functional neuroimaging http://n.neurology.org/cgi/collection/functional_neuroimaging
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2017 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.