Teaching *NeuroImages*: Osteochondroma arising from the clavicle causing ipsilateral Horner syndrome

Diana Y. Wei, MBBS,
MRCP
Mike Bradley, MBChB
Zolt Orosz, MD, PhD
James C. Stevens, MBBS,
PhD

Correspondence to
D. Wei:
dianawei@doctors.org.uk

Download teaching slides:
Neurology.org

A 17-year-old female smoker noticed right eyelid droop for 6 months, with a constant right-sided pressure-like headache with photophobia, phonophobia, and nausea, right-sided lacrimation, and right-sided rhinorrhea. Examination revealed right Horner syndrome. A mass was palpable on the anterior aspect of her neck. Ultrasound of the neck, MRI head, and CT thorax showed a bony mass arising from the right medial clavicular head (figure 1). Histopathology from the excision confirmed an osteochondroma (figure 2). This was removed surgically with partial resolution of symptoms. This case demonstrates a rare cause of Horner syndrome^{1,2} and the importance of thorough imaging of the sympathetic chain.

AUTHOR CONTRIBUTIONS

Diana Wei: wrote the initial manuscript and acquisition of data and review of manuscript. Mike Bradley: interpretation of CT and MRI

scans. Zolt Orosz: interpretation of histopathology slides. James Stevens: case report concept and critical revision of manuscript for intellectual content.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

D. Wei, M. Bradley, and Z. Orosz report no disclosures relevant to the manuscript. J. Stevens served on an advisory board for Grifols UK. Go to Neurology.org for full disclosures.

REFERENCES

1. Simpson JF. Horner's syndrome due to an osteochondroma of the first rib. *Can Med Assoc J* 1948;59:152–155.
2. Zhao CQ, Jiang SD, Jiang LS, Dai LY. Horner Syndrome due to a solitary osteochondroma of C7: a case report and review of the literature. *Spine* 2007;32:E471–E474.

Figure 1 Imaging showing the bony mass

(A) Coronal CT bone window shows the osteochondroma arising from the medial clavicle. (B) MRI STIR axial and (C) coronal demonstrate the osteochondroma in relation to right carotid and subclavian and cartilage cap.

Figure 2 Histopathology from the excision confirmed an osteochondroma

Neurology®

Teaching *NeuroImages*: Osteochondroma arising from the clavicle causing ipsilateral Horner syndrome

Diana Y. Wei, Mike Bradley, Zsolt Orosz, et al.

Neurology 2017;89:e13-e14

DOI 10.1212/WNL.0000000000004084

This information is current as of July 10, 2017

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/89/2/e13.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2017/07/10/WNL.0000000000004084.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/89/2/e13.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology CT http://n.neurology.org/cgi/collection/ct MRI http://n.neurology.org/cgi/collection/mri Pupils http://n.neurology.org/cgi/collection/pupils
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2017 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.