Amyloid polyneuropathy in 2 patients after liver transplantation

Ainoa Ugarte, MD, Josep M. Grau, MD, PhD, and José C. Milisenda, MD

Neurology® 2018;90:38. doi:10.1212/WNL.0000000000004777

Correspondence

Dr. Milisenda

jcmilise@clinic.ub.es

Figure Peripheral nerve biopsies

Transverse sural nerve sections, paraffin embedded. (A) Positive Congo red staining in the endoneurial vessels (star). (B) Positive prealbumin (transthyretin) immunohistochemistry in the endoneurial vessels (arrows).

Domino liver transplantation, in which the liver from a patient with transthyretin-mediated amyloidosis (ATTR) is transplanted into another patient, is a well-established procedure.^{1,2} The figure is from 2 unrelated cases of liver transplant recipients from donors with ATTR who developed clinical polyneuropathy with weakness and paresthesia in lower extremities some years after the liver transplantation. In both cases, EMG showed distal sensory and motor polyneuropathy. Although amyloid neuropathy was the clinical suspicion, a muscle and nerve biopsy was performed in order to rule out other conditions.

Author contributions

Ainoa Ugarte: acquisition of data. Josep M. Grau: critical revision of the manuscript for important intellectual content. José C. Milisenda: study concept and design.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

1. Dixit N, Castano A, Farr MJ, et al. Rapidly progressive transthyretin-mediated amyloidosis in a domino liver transplant recipient of a Ser23Asn donor. *J Clin Neuromuscul Dis* 2016;17:142–145.
2. Adams D, Cauquil C, Labeyrie C. Familial amyloid polyneuropathy. *Curr Opin Neurol* 2017;30:481–489.

Neurology®

Amyloid polyneuropathy in 2 patients after liver transplantation

Ainoa Ugarte, Josep M. Grau and José C. Milisenda

Neurology 2018;90;38

DOI 10.1212/WNL.0000000000004777

This information is current as of December 28, 2017

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/90/1/38.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/90/1/38.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology All Neurotology http://n.neurology.org/cgi/collection/all_neurotology EMG http://n.neurology.org/cgi/collection/emg Peripheral neuropathy http://n.neurology.org/cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2017 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.