Atypical subependymal giant cell astrocytoma and neonatal tuberous sclerosis

Laiz Laura de Godoy, MD, and César Augusto Pinheiro Ferreira Alves, MD

Neurology® 2018;90:570-571. doi:10.1212/WNL.0000000000005181

Correspondence

Dr. de Godoy

laizgodoy@hotmail.com

Figure 1 Brain MRI

Axial MRI of tuberous sclerosis brain findings. Subependymal giant astrocytoma: (A) T1-hyperintense; (B) T2-hypointense; (C) susceptibility-weighted imaging—exuberant venous drainage (arrowheads). Subependymal nodules: (A) T1-hyperintense (black arrows), radial bands (red arrows); and (B) cortical tubers (asterisk).

A 9-day-old girl had, on prenatal ultrasound, brain and cardiac lesions suspicious for tuberous sclerosis. Brain MRI demonstrated a large intraventricular and intraparenchymal mass centered on the foramen of Monro, which had unusual imaging findings: a homogeneous T1-hyperintense and T2-hypointense signal (figure 1, A and B) related to scarce myelination; remarkable venous drainage on susceptibility-weighted imaging (figure 1C); and apparent diffusion coefficient hypointensity, corresponding to blackout T2 effect (figure 2, A and B).

Figure 2 Brain MRI

Subependymal giant astrocytoma: (A) apparent diffusion coefficient hypointensity; (B) diffusion-weighted imaging isointense related to blackout T2 effect; (C) spectroscopy (point resolved spectroscopy echo time 35) study shows choline (Cho) 3.2 ppm and myo-inositol (Myo) 3.5 ppm peaks.

Spectroscopy showed increased choline and myo-Inositol peaks (figure 2C). These findings are consistent with neonatal subependymal giant astrocytoma.^{1,2} Subependymal nodules, cortical tubers, and radial bands were also atypically T1-hyperintense and T2-hypointense (figure 1, A and B).

Author contributions

Laiz Laura de Godoy: study concept and design. César Augusto Pinheiro Ferreira Alves: study concept and design, critical revision of manuscript for intellectual content.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

1. Raju GP, Urion DK, Sahin M. Neonatal subependymal giant cell astrocytoma: new case and review of literature. *Pediatr Neurol* 2007;36:128–131.
2. Baron Y, Barkovich AJ. MR imaging of tuberous sclerosis in neonates and young infants. *AJNR Am J Neuroradiol* 1999;20:907–916.

Disputes & Debates: Rapid online correspondence

The editors encourage comments on recent articles through Disputes & Debates:

Access an article at Neurology.org/N and click on “COMMENT” beneath the article header. Responses will be posted within 3 business days.

Before submitting a comment to Disputes & Debates, remember the following:

- Disputes & Debates is restricted to comments about studies published in *Neurology* within the last eight weeks
 - Read previously posted comments; redundant comments will not be posted
 - Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
 - You can include a maximum of five authors (including yourself)
-

The AAN is Fighting for You!

In the midst of rapid changes in health care policy, the AAN has your back. From actively lobbying members of Congress to meeting with regulators to underscore the value of neurology and your services to your patients, the Academy is forcefully countering any threats to your profession and patient access to care. Learn more at AAN.com/policy-and-guidelines/advocacy, read the bimonthly Capitol Hill Report and monthly *AANnews*® member magazine, and respond to Advocacy Action Alert emails when we invite you to share your voice with Congress.

Neurology®

Atypical subependymal giant cell astrocytoma and neonatal tuberous sclerosis

Laiz Laura de Godoy and César Augusto Pinheiro Ferreira Alves

Neurology 2018;90;570-571

DOI 10.1212/WNL.0000000000005181

This information is current as of March 19, 2018

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/90/12/570.full
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/90/12/570.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): DWI http://n.neurology.org/cgi/collection/dwi MRI http://n.neurology.org/cgi/collection/mri Neonatal http://n.neurology.org/cgi/collection/neonatal Other neurocutaneous disorders http://n.neurology.org/cgi/collection/other_neurocutaneous_disorders
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.