Abstracts

Articles appearing in the February 2018 issue

Diagnostic utility of exome sequencing in the evaluation of neuromuscular disorders

Objective To evaluate the diagnostic yield and workflow of genome-scale sequencing in patients with neuromuscular disorders (NMDs).

Methods We performed exome sequencing in 93 undiagnosed patients with various NMDs for whom a molecular diagnosis was not yet established.

Variants on both targeted and broad diagnostic gene lists were identified. Prior diagnostic tests were extracted from the patients' medical records to evaluate the use of exome sequencing in the context of their prior diagnostic workup.

Results The overall diagnostic yield of exome sequencing in our cohort was 12.9%, with 1 or more pathogenic or likely pathogenic variants identified in a causative gene associated with the patient's disorder. Targeted gene lists had the same diagnostic yield as a broad NMD gene list in patients with clear neuropathy or myopathy phenotypes, but evaluation of a broader set of disease genes was needed for patients with complex NMD phenotypes. Most patients with NMD had undergone prior testing, but only 10/16 (63%) of these procedures, such as muscle biopsy, were informative in pointing to a final molecular diagnosis.

Conclusions Genome-scale sequencing or analysis of a panel of relevant genes used early in the evaluation of patients with NMDs can provide or clarify a diagnosis and minimize invasive testing in many cases.

NPub.org/NG/9016a

Alzheimer risk loci and associated neuropathology in a population-based study (Vantaa 85+)

Objective To test the association of distinct neuropathologic features of Alzheimer disease (AD) with risk loci identified in genome-wide association studies.

Methods Vantaa 85+ is a population-based study that includes 601 participants aged ≥ 85 years, 256 of whom were neuropathologically examined. We analyzed 29 AD risk loci in addition to *APOE* $\epsilon 4$, which was studied separately and used as a covariate. Genotyping was performed using a single nucleotide polymorphism (SNP) array (341 variants) and imputation (6,038 variants). Participants with Consortium to Establish a Registry for Alzheimer Disease (CERAD) (neurotic β -amyloid [$A\beta$] plaques) scores 0 ($n = 65$) vs score M + F ($n = 171$) and Braak (neurofibrillary tangle pathology) stages 0–II ($n = 74$) vs stages IV–VI ($n = 119$) and with capillary $A\beta$ (Cap $A\beta$, $n = 77$) vs without ($n = 179$) were compared. Cerebral amyloid angiopathy (CAA) percentage was analyzed as a continuous variable.

Results Altogether, 24 of the 29 loci were associated (at $p < 0.05$) with 1 or more AD-related neuropathologic features in either SNP array or imputation data. Fifteen loci associated with CERAD score, smallest $p = 0.0002122$, odds ratio (OR) 2.67 (1.58–4.49) at *MEF2C* locus. Fifteen loci associated with Braak stage, smallest $p = 0.004372$, OR 0.31 (0.14–0.69) at *GAB2* locus. Twenty loci associated with CAA, smallest $p = 7.17E-07$, β 14.4 (8.88–20) at *CRI* locus. Fifteen loci associated with Cap $A\beta$, smallest $p = 0.002594$, OR 0.54 (0.37–0.81) at *HLA-DRB1* locus. Certain loci associated with specific neuropathologic features. *CASS4*, *CLU*, and *ZCWPW1* associated only with CAA, while *TREM2* and *HLA-DRB5* associated only with Cap $A\beta$.

Conclusions AD risk loci differ in their association with neuropathologic features, and we show for the first time distinct risk loci for CAA and Cap $A\beta$.

NPub.org/NG/9016b

Most-Read Articles

As of February 7, 2018

Mendelian randomization shows a causal effect of low vitamin D on multiple sclerosis risk

B. Rhead, M. Bäärnhielm, M. Gianfrancesco, et al. 2016;2:e97.

CHCHD10 variant p.(Gly66Val) causes axonal Charcot-Marie-Tooth disease

M. Auranen, E. Ylikallio, M. Shcherbii, et al. 2015;1:e1.

A novel *DYNC1H1* mutation causing spinal muscular atrophy with lower extremity predominance

Q. Niu, X. Wang, M. Shi, Q. Jin. 2015;1:e20.

The Clinical Outcome Study for dysferlinopathy: An international multicenter study

E. Harris, C.L. Bladen, A. Mayhew, et al. 2016;2:e89.

Complete callosal agenesis, pontocerebellar hypoplasia, and axonal neuropathy due to *AMPD2* loss

A.P.L. Marsh, V. Lukic, K. Pope, et al. 2015;1:e16.

Neurology[®]

What's happening in *Neurology*[®] *Genetics*
Neurology 2018;90;735
DOI 10.1212/WNL.0000000000005342

This information is current as of April 16, 2018

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/90/16/735.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.