Abstracts

Articles appearing in the December 2017 issue

Incorporating students into clinic may be associated with both improved clinical productivity and educational value

Background In this study, we aimed to evaluate ambulatory clinic responsibilities that neurology clerkship students perceive as having the highest educational value and to evaluate the association between a student's presence and level of responsibility and a preceptor's clinical and financial productivity during a clinic session.

Methods Physician preceptors ($n = 43$) and medical students ($n = 67$) in the Johns Hopkins Neurology clerkship from 2014 to 2015 were included. Students rated their experience and responsibilities in 291 neurology clinic sessions. Productivity metrics (e.g., relative value units [RVU]/clinic) were collected for each preceptor in the presence and absence of students.

Results A student's rating of a clinic as an effective learning experience increased with each additional patient the student interviewed (odds ratio [OR] 1.89, $p < 0.001$), presented (OR 1.86, $p < 0.001$), or documented (OR 2.00, $p < 0.001$). The mean RVU/session for preceptors also increased based on the number of patients interviewed ($\beta = 2.64$, $p = 0.026$), presented ($\beta = 2.42$, $p = 0.047$), and documented ($\beta = 2.70$, $p = 0.036$) by students. On average, preceptor RVU/session increased by 42% (mean 5.6 ± 1.2 , $p < 0.0001$) when a student was present in clinic compared to sessions without students. In addition, preceptor invoices increased by 35% (mean 2.7 ± 0.6 , $p < 0.0001$) and charges by 39% (mean $\$929 \pm \$210$, $p < 0.0001$) when a student was present in clinic.

Conclusions This observational study suggests a mutual benefit to preceptor clinical productivity and student-perceived educational value when students have active responsibilities in neurology clinics. Despite concerns that students slow down preceptors in clinic, these results suggest that preceptors may have an overall boost in productivity, potentially by performing billable work while students independently see patients.

NPub.org/NCP/9007a

➔ Editorial

NPub.org/NCP/9007b

Practice Current

Our latest survey features a hot topic for all practicing neurologists worldwide: When do you suspect autoimmune encephalitis (AE) and what is the role of antibody testing? There has been growing interest in stressing the importance of clinically grounded guidelines for diagnosis of AE and antibody testing may not be available in diverse settings with varying resources. We welcome readers to join the discussion.

NPub.org/NCP/pc6

The upgoing thumb sign: An interobserver/intraobserver reliability study

Background The upgoing thumb sign as a subtle clinical finding of upper motor neuron involvement has been reported frequently in patients with TIAs and minor strokes. This study was designed to show the method of examination and interpretation and the interobserver/intraobserver reliability.

Methods The thumb sign was elicited in TIA/minor strokes or stroke mimics. After obtaining the participant's permission, the examinations were recorded. Two independent neurologists reviewed all patients for the possibility of an upgoing thumb sign. After 1 hour education about the definition of an upgoing thumb sign to a group of stroke fellows, nurse practitioners, and a research secretary, the videos were reviewed and the participants rated the thumb sign independently. The intraobserver reliability was assessed after 3 months by rating the same videos.

Results The interobserver reliability among 9 raters showed an overall agreement of 0.83 and fixed-marginal kappa of 0.66. The same videos were reviewed by the observers after 3 months with a similar level of agreement (percent of overall agreement 0.84, fixed-marginal kappa 0.66) and a substantial to almost perfect level of intraobserver concordance (mean 0.86; SD 0.08; median 0.90; interquartile range 25–75, 0.8, 0.95).

Conclusions The upgoing thumb sign is a subtle upper limb neurologic finding, with a high level of interobserver and intraobserver reliability. The test is easy to perform and can be interpreted accurately.

NPub.org/NCP/9007c

Neurology®

What's happening in *Neurology*® *Clinical Practice*
Neurology 2018;90;323
DOI 10.1212/WNL.0000000000004956

This information is current as of February 12, 2018

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/90/7/323.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.