

Prestroke physical activity could influence acute stroke severity (part of PAPSIGOT)

Malin Reinholdsson, PT, MS, Annie Palstam, PT, PhD, and Katharina S. Sunnerhagen, MD, PhD, Prof

Cite as: *Neurology*® 2018;91:e1461-e1467. doi:10.1212/WNL.0000000000006354

Correspondence

Dr. Reinholdsson
malin.reinholdsson@
vgregion.se

Study objective

To determine how prestroke physical activity (PA) affects acute stroke severities.

Summary results

Prestroke PA is associated with milder stroke severities.

What is known and what this paper adds

Animal studies have indicated that prestroke PA reduces stroke severities and promotes recovery, but human studies have been inconsistent. This study provides evidence for the benefits of prestroke PA in humans.

Participants and setting

This study reviewed data for 925 patients (45.2% female; mean age, 73.1 years; age range, 20–104 years) who were admitted to Sahlgrenska University Hospital (Gothenburg, Sweden) for first-ever intracerebral hemorrhage or ischemic stroke between November 1, 2014, and April 30, 2016.

Design, size, and duration

This study retrospectively accessed data from the Swedish Stroke Register and a local stroke register called Väststroke. NIH Stroke Scale (NIHSS) scores were used to classify stroke events as mild (NIHSS scores of 0–5) or moderate-to-severe (NIHSS scores ≥ 6). Physical therapists asked the patients about their prestroke PA levels and then assigned Saltin-Grimby PA Level Scale (SGPALS) scores, with scores of 1, 2, and 3–4 indicating prestroke physical inactivity, light prestroke PA, and moderate to high prestroke PA, respectively. Logistic regression was used to identify predictors of stroke severity, and negative binomial regression analyses were used to specifically examine associations between prestroke PA and stroke severity.

Primary outcome measures

The primary outcome was the association between prestroke PA and stroke severity.

Table Predictors of mild stroke severities in the logistic regression model

Predictor of mild stroke severities	B Coefficient (standard error)	OR (95% CI)
Prestroke PA	0.70 (0.18)	2.02 (1.43–2.86)
Age	–0.03 (0.01)	0.97 (0.96–0.99)

Main results and the role of chance

Negative binomial regression showed that prestroke physical inactivity was associated with greater stroke severities relative to light prestroke PA ($p < 0.001$) and moderate prestroke PA ($p < 0.001$). The predictors of mild stroke severities in the logistic regression model were prestroke PA and younger ages ($p < 0.001$ for both), but the model's explanatory value was only 6.8%.

Bias, confounding, and other reasons for caution

Some data were missing for some participants. SGPALS scores were assigned based on self-reported prestroke PA, which might have introduced recall bias.

Generalizability to other populations

The large patient sample from a country with freely accessible healthcare favors the generalizability of this study's results.

Study funding/potential competing interests

This study was funded by various foundations and Swedish national and local government agencies. The authors report no competing interests. Go to Neurology.org/N for full disclosures.

A draft of the short-form article was written by M. Dalefield, a writer with Editage, a division of Cactus Communications. The authors of the full-length article and the journal editors edited and approved the final version.

Neurology[®]

Prestroke physical activity could influence acute stroke severity (part of PAPSIGOT)

Malin Reinholdsson, Annie Palstam and Katharina S. Sunnerhagen

Neurology 2018;91:e1461-e1467 Published Online before print September 19, 2018

DOI 10.1212/WNL.0000000000006354

This information is current as of September 19, 2018

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/91/16/e1461.full
References	This article cites 34 articles, 6 of which you can access for free at: http://n.neurology.org/content/91/16/e1461.full#ref-list-1
Citations	This article has been cited by 4 HighWire-hosted articles: http://n.neurology.org/content/91/16/e1461.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Cerebrovascular disease/Stroke http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke All Rehabilitation http://n.neurology.org/cgi/collection/all_rehabilitation Natural history studies (prognosis) http://n.neurology.org/cgi/collection/natural_history_studies_prognosis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

