Teaching NeuroImages: Japanese encephalitis

Bingjun Zhang, MD, Siyuan Liao, MD, Yu Yang, PhD, and Zhengqi Lu, PhD

Neurology® 2018;91:e2031-e2032. doi:10.1212/WNL.0000000000006553

Correspondence

Dr. Lu
lzqzssy@gmail.com

Figure MRI findings in Japanese encephalitis

(A–C) MRI brain axial T2 fluid-attenuated inversion recovery: symmetric hyperintensity in the thalami and substantia nigra. (D–F) MRI brain axial T1+ gadolinium: no contrast enhancement. (G–I) MRI brain axial T2 fluid-attenuated inversion recovery: The lesions had improved substantially after 3 months supportive care.

A 20-year-old woman presented with 3 days' history of fever and deteriorating consciousness. Physical examination showed positive Kerning sign. Brain MRI revealed hyperintensity involving bilateral thalami and substantia nigra (figure). Although CSF tests

MORE ONLINE

→Teaching slides

links.lww.com/WNL/A742

From the Department of Neurology, the Third Affiliated Hospital of Sun Yat-sen University, Guangzhou, China.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Copyright © 2018 American Academy of Neurology

Copyright © 2018 American Academy of Neurology. Unauthorized reproduction of this article is prohibited.

e2031

for white blood cells, protein, glucose, chlorine, and TORCH antibodies were normal, Japanese encephalitis (JE) virus immunoglobulin M antibody was positive. Therefore, a diagnosis of JE was made. The patient recovered from coma, and brain lesions were disappearing after 3 months supportive care (figure). As a common human viral encephalitis in the world, JE is usually very severe with high case-fatality rate.¹ Symmetric thalami and substantia nigra involvement are characteristic in JE.²

Author contributions

B. Zhang: study concept and design, acquisition and analysis of data, preparation of manuscript including figures. S. Liao: data analysis and interpretation. Y. Yang: acquisition and

preparation of data. Z. Lu: study concept and design, data acquisition and interpretation, critical revision.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

1. Misra UK, Kalita J. Overview: Japanese encephalitis. *Prog Neurobiol* 2010;91:108–120.
2. Dung NM, Turtle L, Chong WK, et al. An evaluation of the usefulness of neuroimaging for the diagnosis of Japanese encephalitis. *J Neurol* 2009;256:2052–2060.

Neurology®

Teaching NeuroImages: Japanese encephalitis

Bingjun Zhang, Siyuan Liao, Yu Yang, et al.

Neurology 2018;91:e2031-e2032

DOI 10.1212/WNL.0000000000006553

This information is current as of November 19, 2018

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/91/21/e2031.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/91/21/e2031.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Meningitis http://n.neurology.org/cgi/collection/meningitis MRI http://n.neurology.org/cgi/collection/mri Viral infections http://n.neurology.org/cgi/collection/viral_infections
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.