Abstracts

Articles appearing in the April 2018 issue

Multiple sclerosis management and reproductive changes: A guide for general neurologists

Purpose of review Multiple sclerosis (MS) disease activity and symptoms are tied to hormonal changes. This review explains the current standard of care in MS at various stages of a woman's reproductive life and helps neurologists answer patients' most common questions surrounding MS care and fertility, pregnancy, and menopause.

Recent findings Recent work has focused on MS risk and exacerbation with variables related to reproductive health. Management of disease-modifying therapies prenatally and postnatally is also a focus.

Summary This review is a concise, practical guide for general neurologists caring for women with MS. MS is a disease that requires adaptation of management as a woman moves through reproductive stages. With proper planning and management, pregnancy is safe for women with MS. We describe the current standard of care based on trials, when available, and on expert opinion.

NPub.org/NCP/9104a

Apathy and functional disability in behavioral variant frontotemporal dementia

Background Behavioral variant frontotemporal dementia (bvFTD) has profound consequences on patients and their families. In this multicenter study, we investigated the contribution of cognitive and neuropsychiatric factors to everyday function at different levels of overall functional impairment.

Methods In a retrospective cross-sectional study, 109 patients with bvFTD from 4 specialist frontotemporal dementia centers (Australia, England, India, and Brazil) were included. The measures administered evaluated everyday function (Disability Assessment for Dementia [DAD]), dementia staging (Clinical Dementia Rating), general cognition (Addenbrooke's Cognitive Examination-revised [ACE-R]), and neuropsychiatric symptoms (Neuropsychiatric Inventory [NPI]). Patients were then subdivided according to functional impairment on the DAD into mild, moderate, severe, and very severe subgroups. Three separate multiple linear regression analyses were run, where (1) total DAD, (2) basic activities of daily living (BADL), and (3) instrumental activities of daily living (IADL) scores were dependent variables; ACE-R total score and selected NPI domains (agitation/aggression, euphoria, apathy, disinhibition, irritability, aberrant motor behavior) were used as independent variables. Age, sex, education, and country of origin were controlled for in the analyses.

Results Cognitive deficits were similar across the mild, moderate, and severe subgroups but significantly worse in the very severe subgroup. NPI domain scores (agitation/aggression, euphoria, apathy, disinhibition, irritability, aberrant motor behavior) did not differ across the DAD subgroups. In the multiple regression analyses, a model including ACE-R and NPI apathy explained 32.5% of the variance for total DAD scores. For IADL, 35.6% of the variance was explained by the ACE-R only. No model emerged for BADL scores.

Conclusions Cognitive deficits and apathy are key contributors to functional disability in bvFTD but factors underlying impairment in BADLs remain unclear. Treatments targeting reduction of disability need to address apathy and cognitive impairment to ensure greater efficacy, especially in regards to IADLs.

NPub.org/NCP/9104b

Practice Current

Our survey on the topic "When do you suspect autoimmune encephalitis and what is the role of antibody testing?" has received over 1,000 responses from over 80 countries. Explore this topic and others on our redesigned website: compare your practice with peers and see survey results displayed on an interactive world map.

NPub.org/NCP/pc6

Neurology[®]

What's happening in *Neurology*[®] *Clinical Practice*
Neurology 2018;91;171
DOI 10.1212/WNL.0000000000005880

This information is current as of July 23, 2018

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/91/4/171.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.