Teaching NeuroImages: Cerebral liposuction

Kevin T. Gobeske, MD, PhD, MPH, Simon Maltais, MD, PhD, Harry J. Cloft, MD, PhD, and Eelco F.M. Wijdicks, MD, PhD

Neurology® 2018;91:e692-e693. doi:10.1212/WNL.0000000000005999

Correspondence

Dr. Gobeske
Gobeske.Kevin@mayo.edu

MORE ONLINE

→Teaching slides

links.lww.com/WNL/A629

A 61-year-old woman showed left hemiplegia and neglect while recovering 6 hours after aortic valve repair. CT showed a hypodense middle cerebral artery segment. CT angiography and CT perfusion (CTP) revealed a carotid terminus occlusion and a large penumbra around a small ischemic core. This prompted cerebral angiography, and embolectomy by Sofia catheter aspiration removed a 1.5-cm piece of adipose tissue, presumably pericardial fat introduced during aortotomy (figure). The neglect, hemianopia, and hemiparesis improved thereafter. This demonstrates a successful fat embolus aspiration to treat stroke after cardiac surgery, and which should be considered if deficits are recognized early and CTP shows significant salvageable tissue.^{1,2}

Author contributions

Kevin Gobeske: study concept and design, acquisition of data, analysis, interpretation, and writing. Simon Maltais: acquisition of data, analysis and interpretation. Harry Cloft: acquisition of data. Eelco Wijdicks: study concept and design, acquisition of data, analysis, interpretation, and writing.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

1. Lee TC, Bartlett ES, Fox AJ, Symons SP. The hypodense artery sign. *AJNR Am J Neuroradiol* 2005;26:2027–2029.
2. Abend NS, Levine JM. Hypodense middle cerebral artery with fat embolus. *Neurocrit Care* 2007;6:147–148.

(A) CT with hypodense middle cerebral artery (arrow) and limited early ischemia (arrowheads). (B) Cerebral angiogram with ICA occlusion (arrow). (C) CT perfusion (CTP) blood flow scan shows large ischemic penumbra. (D) CTP blood volume scan shows small ischemic core. (E) Embolectomy gross pathology and (F) 40× hematoxylin & eosin stain photomicrograph demonstrate adipose tissue.

Neurology®

Teaching NeuroImages: Cerebral liposuction

Kevin T. Gobeske, Simon Maltais, Harry J. Cloft, et al.

Neurology 2018;91:e692-e693

DOI 10.1212/WNL.0000000000005999

This information is current as of August 13, 2018

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/91/7/e692.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/91/7/e692.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Imaging http://n.neurology.org/cgi/collection/all_imaging Critical care http://n.neurology.org/cgi/collection/critical_care CT http://n.neurology.org/cgi/collection/ct Embolism http://n.neurology.org/cgi/collection/embolism Infarction http://n.neurology.org/cgi/collection/infarction
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.