Clinical Reasoning: A 15-year-old boy with bilateral wrist pain in the setting of weight loss

K.H. Vincent Lau, MD, William S. David, MD, PhD, and Reza Sadjadi, MD

Neurology® 2019;92:486-492. doi:10.1212/WNL.0000000000007034

Correspondence

Dr. Sadjadi
rseyedsadjadi@
mgh.harvard.edu

Section 1

A 15-year-old previously healthy boy was referred for further evaluation of wrist pain. Right and left wrist pain started 2 years ago and got significantly worse in the setting of intentional 25-pound weight loss over 6 months. He also had numbness in the 4th and 5th digits of his left hand for the last 3 months. Neurologic examination was notable for weakness of both thumb abduction, finger abduction, and flexion of 4th and 5th digits on the left. There was decreased sensation to pinprick in palms, as well as trace deep tendon reflexes at biceps, triceps, and brachioradialis.

Questions for consideration:

1. What is the localization of the patient's deficits?
2. What is the best next diagnostic step?

GO TO SECTION 2

From the Neuromuscular Division, Massachusetts General Hospital, Boston.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Section 2

The patient's deficits localize to bilateral median nerves (thumb abduction) and bilateral ulnar nerves (finger abduction and flexion of left 4th and 5th digits). To better localize the neuropathies—for example, to determine if there is evidence of

focal slowing at the carpal tunnel for the median nerves—the best next diagnostic step is nerve conduction studies (NCS). These were performed and are summarized in the table.

Question for consideration:

1. What is the interpretation of these NCS?

Table Nerve conduction studies

Nerve/sites	Distance, cm	Peak latency, ms	Amplitude, μ V	Temp., $^{\circ}$ C	
Sensory					
R median—Digit II					
Digit II	13	4.4 ^a	7.4 ^a	29.4	
Palm	8	3.2	46.3	29.3	
L median—Digit II					
Digit II	13	3.3 ^a	12.0	33.4	
Palm	8	2.4	38.8	33.4	
R ulnar—Digit V					
Digit V	11	3.2	15.2	31.7	
Palm	8	2.4	39.8	30.8	
L ulnar—Digit V					
Digit V	11	2.7	14.2	32.6	
Palm	8	2.1	40.0	32.4	
R radial—snuff box					
Forearm	10	2.9	32.8	28.3	
L radial—snuff box					
Forearm	10	2.4	35.7	32.7	
R sural—lateral malleolus					
Calf	14	4.8 ^a	19.7	30.8	
L sural— lateral malleolus					
Calf	14	4.5 ^a	12.4	30.6	
R superficial peroneal					
Lateral leg	14	4.0	7.8	30.6	
L superficial peroneal					
Lateral leg	14	4.3	8.4	30.1	
Nerve/sites	Distance, cm	Latency, ms	Amplitude, mV	Velocity, m/s	Temp., $^{\circ}$ C
Motor					
R median—APB					
Wrist	6	5.7 ^a	13.5		30.3
Elbow	23.5	10.4	13.3	50.1	30.1

Continued

Table Nerve conduction studies (*continued*)

Nerve/sites	Distance, cm	Latency, ms	Amplitude, mV	Velocity, m/s	Temp., °C
L median—APB					
Wrist	6	4.9 ^a	9.8		33.8
Elbow	22.5	9.1	9.8	54.0	33.8
R ulnar—ADM					
Wrist	6	3.3	14.5		28.4
B. elbow	22	8.1	14.3	45.9	28.4
A. elbow	10	11.4	13.4	30.0 ^a	28.3
L ulnar—ADM					
Wrist	6	2.4	11.0		33.4
B. elbow	21.5	6.5	10.9	52.3	33.5
A. elbow	10	9.7	9.4	31.5 ^a	33.8
R ulnar—FDI					
Wrist		3.6	10.8		32.4
B. elbow	22	8.4	9.7	45.9	32.4
A. elbow	10	11.7	8.0	30.5 ^a	32.1
L ulnar—FDI					
Wrist		3.8	9.2		32.2
B. elbow	21.5	7.8	9.2	52.9	32.4
A. elbow	10	10.9	6.5	32.0 ^a	32.9
R peroneal—EDB					
Ankle	9	5.6	8.4		30.8
Below fibular head	31	14.7	6.6	34.0 ^a	31
Above fibular head	9	16.7	6.3	46.7	30.8
L peroneal—EDB					
Ankle	9	5.9	7.9		30.2
Below fibular head	32	15.1	6.6	35.1 ^a	30.2
Above fibular head	9	17.0	6.2	45.5	30.4
R tibial—AH					
Ankle	10	6.0	11.2		31.3
Popliteal fossa	39.5	16.9	8.3	36.1 ^a	31.1
L tibial—AH					
Ankle	10	5.4	8.0		30.6
Popliteal fossa	36	15.4	6.7	36.2 ^a	30.6

Abbreviations: A. elbow = above elbow; ADM = abductor digiti minimi; AH = abductor hallucis; APB = abductor pollicis brevis; B. elbow = below elbow; EDB = extensor digitorum brevis; FDI = first dorsal interosseous.

^a Abnormal values.

GO TO SECTION 3

Section 3

NCS demonstrated that bilateral median sensory nerve action potentials (SNAP) had prolonged distal latencies, with decreased amplitude on the right and normal amplitude on the left. Both median compound motor action potentials (CMAPs) had prolonged distal latencies, and bilateral ulnar CMAPs recording from the adductor digiti minimi and first dorsal interossei had focal slowing of conduction velocities across the elbows (i.e., ulnar grooves). Both sural SNAPs had prolonged peak latencies and bilateral peroneal and tibial CMAPs

had slowed conduction velocities. Taken together, there was electrodiagnostic evidence for bilateral moderate median neuropathies at the wrist, i.e., carpal tunnel syndrome, and bilateral ulnar neuropathies localized to the elbows, superimposed on a generalized, demyelinating, sensorimotor polyneuropathy.

A month prior to the patient's referral visit, he had an ultrasound study of bilateral median and ulnar nerves (figure).

Question for consideration:

1. What is the interpretation of these ultrasound images?

Figure Ultrasound images of bilateral median and ulnar nerves

Ultrasound images demonstrate cross-sectional areas of (A) left ulnar nerve, (B) right ulnar nerve, (C) left median nerve at forearm (left) and wrist (right), and (D) right median nerve at wrist (left) and forearm (right).

GO TO SECTION 4

Section 4

Ultrasound demonstrated abnormal ratios of wrist and mid-forearm cross-sectional areas (CSA) of bilateral median nerves (2.29 on right, 1.66 on left; normal <1.4) and increased CSA of bilateral ulnar nerves (13.5 mm² on left, 21.1 mm² on right; normal <6.5 mm²). In nerve ultrasound, increased nerve CSA corresponds with swelling, which can be seen in any focal neuropathy; for example, the median nerve at the wrist in carpal tunnel syndrome. A nerve segment is considered abnormally large when its CSA is increased in absolute terms or

relative to a distant, presumably normal segment.¹ In this case, increased ratio of median nerve CSA at the wrists relative to the mid-forearms and increased absolute ulnar nerve CSA at the elbows suggest neuropathies localized to these compressible sites.

Questions for consideration:

1. What are the differential diagnoses for this patient's presentation?
2. What is the most likely diagnosis?
3. What test would confirm the diagnosis?

GO TO SECTION 5

Section 5

The presentation of multiple focal mononeuropathies is suggestive of an underlying systemic process, as it would be highly unlikely for multiple neuropathies to occur coincidentally in an otherwise healthy individual. It is important to note that this presentation differs from that of the more common generalized polyneuropathy, for which nerve abnormalities are relatively uniform and not clearly localizable as in this case; this type of polyneuropathy can be seen in acquired disorders such as vitamin B₁₂ deficiency or inherited disorders such as Charcot-Marie-Tooth (CMT) disease.

The differential diagnosis for multiple focal neuropathies is narrow compared to generalized polyneuropathy. In the general population, these include diabetes, connective tissue disorders such as systemic lupus erythematosus, and vasculitic disorders such as polyarteritis nodosa causing mononeuritis multiplex. Less common diagnoses include amyloidosis and leprosy. When multiple focal neuropathies are found at common compressive sites as in bilateral carpal tunnels and ulnar grooves as in this case, one specific diagnosis to consider is hereditary neuropathy with liability to pressure palsy (HNPP). This diagnosis is especially likely in a pediatric patient, in whom the aforementioned disorders associated with multiple focal neuropathies are much less likely compared to for an adult. While other systemic disorders can increase the risk of compressive neuropathy—for example, carpal tunnel syndrome in amyloidosis—multiple compressive neuropathies in different limbs and distributions is highly suggestive of HNPP, which is the most likely diagnosis in this case.

In our patient, targeted genetic testing for HNPP demonstrated heterozygosity for deletion of all 5 exons of the peripheral myelin protein-22 (*PMP22*) gene. He was diagnosed with HNPP with symptoms that were exacerbated by weight loss.

HNPP is a rare autosomal dominant peripheral neuropathy, with prevalence estimated between 0.84 and 16 per 100,000. Interestingly, a recent study in which next-generation sequencing–based copy number variation analysis was performed on asymptomatic newborns found a genetic prevalence characterized by *PMP22* deletion of 58.9 per 100,000, which is unsurprising given the combination of underdiagnosis and likely incomplete penetrance of the disease.² Age at onset of first HNPP symptoms is typically in the second or third decade, with range from birth to the eighth decade. Given the rarity of the disorder, the epidemiology of HNPP in children is not well-established, although it has been suggested that clinical suspicion should be high in children with multifocal neuropathy even without a family history (in the largest case series on pediatric HNPP to date, only 25% had affected family members).³

HNPP is characterized by recurrent peripheral nerve injury leading to weakness or sensory loss precipitated by minor compression or trauma. The weight loss noted in this case is

a well-known risk factor for compression neuropathies in general, the mechanism thought to be multifactorial, including reduction of protective subcutaneous tissue and metabolic changes.⁴ The molecular basis of HNPP pathophysiology is related to interactions of the *PMP22* protein with proteins that regulate myelin junctions.⁵

In HNPP, neurologic examination demonstrates weakness and sensory loss in the distribution of affected nerves, with diffusely reduced deep tendon reflexes in some cases. Electrophysiologic studies demonstrate decreased motor nerve conduction velocities, prolonged distal motor latencies predominantly at sites of common nerve entrapment, and abnormal sensory nerve action potentials even in clinically unaffected nerves.⁶ A recent study suggests that diagnostic workup for suspected HNPP should include a complete nerve conduction study with a minimum of bilateral median, ulnar, and peroneal nerves independent of symptomatology, due to typical generalization of nerve involvement.⁷ Sonographic evaluation may be helpful in cases in which electrodiagnostic studies and genetic testing are unavailable or not completely consistent with HNPP,⁸ likely unnecessary in this case as electrodiagnostic studies were highly suggestive. In terms of genetics, a contiguous gene deletion of chromosome 17p11.2 including *PMP22* is found in approximately 80% of affected patients. The remainder have a pathogenic variant in *PMP22*, the specific mutation of which can cause a continuum of severity, ranging from a minimally symptomatic presentation to one with persistent and progressive deficits approximating a CMT presentation.⁵

The management of HNPP is generally conservative, and patients are advised to avoid external compression and minor trauma. However, recent case reports have suggested benefit with steroids (demonstrated in one patient with improvement in motor function associated with radial nerve and another associated with peroneal nerve)⁹ and IV immunoglobulins (demonstrated in one patient with improved lower extremity pain).¹⁰ These reports suggest that inflammation may be important in the pathophysiology of HNPP, especially in cases in which pain is a major symptom.

Our patient avoided compression at the wrists by using dictation software for school to limit typing time. He also participated in occupational therapy. Two months later, repeat examination demonstrated slight improvement in bilateral thumb abduction strength, and repeat NCS demonstrated slight improvement in amplitudes of bilateral median CMAPs and SNAPs. He was advised to maintain his current weight and to continue avoiding nerve compression.

Author contributions

Dr. Lau drafted the initial manuscript, revised the manuscript, and was involved in the clinical care of the patient. Dr. Sadjadi revised the manuscript and was involved in the clinical care of the patient. Dr. David was involved in the clinical care of the patient.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](#) for full disclosures.

References

1. Suk JJ, Walker FO, Cartwright MS. Ultrasonography of peripheral nerves. *Curr Neurol Neurosci Rep* 2013;13:328.
2. Park JE, Noh SJ, Oh M, Cho DY, Kim SY, Ki CS. Frequency of hereditary neuropathy with liability to pressure palsies (HNPP) due to 17p11.2 deletion in a Korean newborn population. *Orphanet J Rare Dis* 2018;13:40.
3. Chrestian N, McMillan H, Poulin C, Campbell C, Vajsa J. Hereditary neuropathy with liability to pressure palsies in childhood: case series and literature update. *Neuromuscul Disord* 2015;25:693–698.
4. Meylaerts L, Cardinaels E, Vandevenne J, et al. Peroneal neuropathy after weight loss: a high-resolution ultrasonographic characterization of the common peroneal nerve. *Skeletal Radiol* 2011;40:1557–1562.
5. Taioli F, Cabrini I, Cavallaro T, Adler M, Fabrizi GM. Inherited demyelinating neuropathies with micromutations of peripheral myelin protein 22 gene. *Brain* 2011;134:608–617.
6. Verhagen WI, Gabreëls-Festen AA, van Wensen PJ, et al. Hereditary neuropathy with liability to pressure palsies: a clinical, electroneurophysiological and morphological study. *J Neurol Sci* 1993;116:176–184.
7. Robert-Varvat F, Jousserand G, Bouhour F, et al. Hereditary neuropathy with liability to pressure palsy in patients under 30 years old: neurophysiological data and proposed electrodiagnostic criteria. *Muscle Nerve* 2018;57:217–221.
8. Kim SH, Yang SN, Yoon JS, Park BJ. Sonographic evaluation of the peripheral nerves in hereditary neuropathy with liability to pressure palsies: a case report. *Ann Rehabil Med* 2014;38:109–115.
9. Heng HS, Tang SS, Goyal S, Wraige EA, Lim MJ. Beneficial use of steroids in hereditary neuropathy with liability to pressure palsy. *Dev Med Child Neurol* 2012;54:183–186.
10. Vrinten C, Gu X, Weinreich SS, et al. An n-of-one RCT for intravenous immunoglobulin G for inflammation in hereditary neuropathy with liability to pressure palsy (HNPP). *J Neurol Neurosurg Psychiatry* 2016;87:790–791.

Disputes & Debates: Rapid online correspondence

The editors encourage comments on recent articles through Disputes & Debates:

Access an article at [Neurology.org/N](#) and click on “COMMENT” beneath the article header. Responses will be posted within 3 business days.

Before submitting a comment to Disputes & Debates, remember the following:

- Disputes & Debates is restricted to comments about studies published in *Neurology* within the last eight weeks
 - Read previously posted comments; redundant comments will not be posted
 - Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
 - You can include a maximum of five authors (including yourself)
-

Visit the *Neurology*® Resident & Fellow Website

Click on Residents & Fellows tab at [Neurology.org](#).

Now offering:

- *Neurology*® Resident & Fellow Editorial team information
- “Search by subcategory” option
- E-pearl of the Week
- RSS Feeds
- Direct links to Continuum®, Career Planning, and AAN Resident & Fellow pages
- Recently published Resident & Fellow articles
- Podcast descriptions

Find *Neurology*® Residents & Fellows Section on Facebook: <http://tinyurl.com/o8ahsys>

Follow *Neurology*® on Twitter: <http://twitter.com/GreenJournal>

Neurology®

Clinical Reasoning: A 15-year-old boy with bilateral wrist pain in the setting of weight loss

K.H. Vincent Lau, William S. David and Reza Sadjadi

Neurology 2019;92;486-492

DOI 10.1212/WNL.00000000000007034

This information is current as of March 4, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/92/10/486.full
References	This article cites 10 articles, 1 of which you can access for free at: http://n.neurology.org/content/92/10/486.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Pediatric http://n.neurology.org/cgi/collection/all_pediatric EMG http://n.neurology.org/cgi/collection/emg Peripheral neuropathy http://n.neurology.org/cgi/collection/peripheral_neuropathy Ultrasound http://n.neurology.org/cgi/collection/ultrasound
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.