Muscle and not neuronal biomarkers correlate with severity in spinal and bulbar muscular atrophy

Vittoria Lombardi, PhD, Giorgia Querin, MD,* Oliver J. Ziff, MD, MRCP,* Luca Zampieri,* Ilaria Martinelli, MD, Carolin Heller, Martha Foiani, Cinzia Bertolin, PhD, Ching-Hua Lu, MD, PhD, Bilal Malik, PhD, Kezia Allen, Carlo Rinaldi, MD, PhD, Henrik Zetterberg, PhD, Amanda Heslegrave, PhD, Linda Greensmith, PhD, Michael Hanna, FRCP, Gianni Soraru, MD, PhD,‡ Andrea Malaspina, MD, PhD, FRCP,‡ and Pietro Fratta, MD, PhD‡

Correspondence

Dr. Fratta
p.fratta@ucl.ac.uk
or Dr. Malaspina
a.malaspina@qmul.ac.uk

Neurology® 2019;92:e1205-e1211. doi:10.1212/WNL.0000000000007097

Abstract

Objective

To determine whether blood biomarkers of neuronal damage (neurofilament light chain [NfL]), muscle damage (creatine kinase [CK]), and muscle mass (creatinine) are altered in spinal and bulbar muscular atrophy (SBMA) and can be used as biomarkers for disease severity.

Methods

In this multicenter longitudinal prospective study, plasma and serum were collected from 2 cohorts of patients with SBMA in London, United Kingdom (n = 50), and Padova, Italy (n = 43), along with disease (amyotrophic lateral sclerosis [ALS]) and healthy controls, and levels of plasma and serum NfL, CK, and creatinine were measured. Disease severity was assessed by the SBMA Functional Rating Scale and the Adult Myopathy Assessment Tool at baseline and 12 and 24 months.

Results

Blood NfL concentrations were increased in ALS samples, but were unchanged in both SBMA cohorts, were stable after 12 and 24 months, and were not correlated with clinical severity. Normal NfL levels were also found in a well-established mouse model of SBMA. Conversely, CK concentrations were significantly raised in SBMA compared with ALS samples, and were not correlated to the clinical measures. Creatinine concentrations were significantly reduced in SBMA, and strongly and significantly correlated with disease severity.

Conclusions

While muscle damage and muscle mass biomarkers are abnormal in SBMA, axonal damage markers are unchanged, highlighting the relevant primary role of skeletal muscle in disease pathogenesis. Creatinine, but not CK, correlated with disease severity, confirming its role as a valuable biomarker in SBMA.

MORE ONLINE

CME Course

NPub.org/cmelist

*These authors contributed equally to this work.

‡These authors contributed equally to this work.

From the Institute of Neurology (V.L., O.J.Z., L.Z., C.H., M.F., C.-H.L., B.M., H.Z., A.H., L.G., M.H., P.F.), University College London Institute of Neurology, Queen Square, London; Blizzard Institute (V.L., A.M.), Queen Mary, University of London, UK; Department of Neurosciences (G.Q., I.M., C.B., G.S.), University of Padova, Italy; Department of Neurology (C.-H.L.), China Medical University Hospital, Taiwan; Basildon Hospital (K.A.), UK; Department of Physiology, Anatomy and Genetics (C.R.), University of Oxford; UK Dementia Research Institute at UCL (H.Z., A.H.), London, UK; Clinical Neurochemistry Laboratory (H.Z.), Sahlgrenska University Hospital; and the Department of Psychiatry and Neurochemistry (H.Z.), Institute of Neuroscience and Physiology, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by Medical Research Council UK.

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Glossary

ALS = amyotrophic lateral sclerosis; **ALSFRS** = Amyotrophic Lateral Sclerosis Functional Rating Scale; **AMAT** = Adult Myopathy Assessment Tool; **AR** = androgen receptor; **CK** = creatine kinase; **KD** = Kennedy disease; **LMN** = lower motor neuron; **NfL** = neurofilament light chain; **PRL** = progression rate to last visit; **SBG** = streptavidin- β -galactosidase; **SBMA** = spinal and bulbar muscular atrophy; **SBMAFRS** = Spinal and Bulbar Muscular Atrophy Functional Rating Scale.

Spinal and bulbar muscular atrophy (SBMA), also known as Kennedy disease (KD), is a disabling adult onset neuromuscular disorder that affects men and is primarily characterized by slowly progressive weakness and wasting of bulbar and limb muscles.^{1,2} SBMA is caused by the expansion of a CAG repeat in the androgen receptor (AR) gene and is characterized by muscle denervation and loss of lower motor neurons in the spinal cord and the brainstem. Signs of primary skeletal muscle damage, such as muscle fiber splitting, fiber degeneration, and centralized nuclei, are also present in SBMA, and recent work showing that silencing of the mutation in muscle is able to rescue the disease phenotype in disease models has further highlighted the relevance of a primary myopathic component in this disorder.^{3–9}

No therapy is currently available for SBMA and an important limitation to trials for promising therapeutic strategies has been the lack of effective outcome measures. Biomarkers to measure disease progression and therapeutic responses are therefore strongly needed.

Neurofilament light chain (NfL) has been found to be increased in serum and plasma of numerous neurologic conditions, including amyotrophic lateral sclerosis (ALS) and inherited peripheral neuropathies, and to be a promising tool to monitor disease progression.^{10,11} We here use the currently most sensitive methodology to measure NfL in patients with SBMA and in a rodent model of disease,¹² and compare this marker of neuronal damage with measures of muscle damage and loss.

Methods

Standard protocol approvals, registrations, and patient consents

Plasma and serum were prospectively collected with informed consent from 2 cohorts of patients with a genetically confirmed diagnosis of SBMA attending the KD clinic at the National Hospital for Neurology in London, United Kingdom ($n = 50$), and at the University Hospital in Padova, Italy ($n = 43$). Samples from patients with ALS ($n = 53$) and healthy relatives of patients ($n = 73$) were collected in neuromuscular clinics in the same centers.

Approvals were obtained from the East London and the City Research Ethics Committee (09/H0703/27). Written informed consent was obtained from all participants in the study.

Patient evaluation

All patients were recruited in neuromuscular clinics at the National Hospital for Neurology in London, United Kingdom, and at the University Hospital in Padova, Italy. All participants with SBMA carried a pathogenic expansion of the CAG trinucleotide repeat (>38 repeats) in the AR gene. ALS cases all were recruited according to El Escorial probable and definite ALS criteria. Controls were recruited in both centers in order to reduce sources of bias. Healthy controls were excluded if they had coexistent neurologic disease as determined by a symptom and medical history-based questionnaire.

SBMA and ALS disease severity and progression were evaluated using functional rating scale scores (Spinal and Bulbar Muscular Atrophy Functional Rating Scale [SBMAFRS], Amyotrophic Lateral Sclerosis Functional Rating Scale [ALSFRS]), and patients with SBMA were further evaluated with the Adult Myopathy Assessment Tool (AMAT).^{13–15}

Progression rate in patients with ALS was calculated as progression rate to last visit (PRL). This was the ALSFRS–revised (range 1–48, with lower scores corresponding to higher level of neurologic impairment) approximated to 48 at onset minus the score at last visit divided per disease duration expressed in month (ALS-Fast = $PRL >1$; ALS-Slow = $PRL <0.5$).

Sample collection and processing

Samples were processed, stored, and analyzed as previously described.^{10,11} Blood samples from all participants were collected into EDTA-containing tubes and centrifuged at 20°C at 3,500 rpm for 10 minutes within 1 hour. Repeat samples were taken after 1 and 2 years, when available. Plasma and serum were aliquoted and stored at -80°C .

Mouse samples

All experimental procedures were carried out under license from the UK Home Office (Scientific Procedures Act 1986), and following approval by the Ethical Review Panel of UCL Institute of Neurology. Yeast artificial chromosome (YAC) AR100 mice, which express at endogenous levels the human androgen receptor with an expanded CAG repeat,¹² were bred and maintained at UCL Institute of Neurology Biological Services. Heterozygote males carrying the androgen receptor with 100 polyglutamine repeats (pathogenic AR100 mice) were mated with wild-type C57BL/6J females. Only male mice were used in this study and male age-matched wild-type littermates were used as controls. Mouse plasma was obtained from male AR100 mice and controls at 18 months of age, as previously described.^{12,16}

NfL, creatine kinase (CK), and creatinine measurements

The quantitative determination of NfL in human plasma (UK cohort) and serum (Italian cohort) was undertaken by single molecule array (Simoa) technology using a digital immunoassay Simoa HD-1 Analyzer (Quanterix; Lexington, MA) using a commercially available NfL kit from the same vendor. Plasma samples from KD, ALS-Fast, ALS-Slow, and healthy patients, along with KD/WT mice, were equally distributed on each plate and measured in duplicate. Each plate contained a target-specific calibrator (500–0.686 pg/mL) and 2 quality controls (10 and 200 pg/mL).

In the first step, anti-NfL-antibody-coated paramagnetic capture beads, sample, and biotinylated detector antibody are combined. NfL molecules present in the sample are captured by the anti-NfL-antibody-coated capture beads and labeled with biotinylated detector antibodies. After washing, a conjugate of streptavidin- β -galactosidase (SBG) is mixed with the capture beads. SBG binds to the biotinylated detector antibodies, resulting in enzyme labeling of captured NfL. CK and creatinine were measured as part of routine clinical testing.

Statistical analysis

Statistical analysis was performed using GraphPad Prism 5.0 (GraphPad; La Jolla, CA). Patient NfL, CK, and creatinine concentrations were compared using analysis of variance and Bonferroni correction for multiple testing. Mouse NfL levels were compared using Mann-Whitney test. Correlations of CK and creatinine levels with SBMAFRS and AMAT scores were assessed using Spearman correlation coefficients. We included all available samples for this study, given the rarity of the disease and the unknown variability and changes of NfL in SBMA prior to this work. Missing data were omitted from analysis—all numbers for each analysis are reported in the table.

Data availability

Anonymized data not published within this article will be made available by request from any qualified investigator.

Results

Cohorts

The study included 93 patients with SBMA, 53 patients with ALS, and 73 healthy controls. Participant characteristics are summarized in the table.

NfL levels are unchanged in 2 cohorts of patients with SBMA

We measured the plasma levels of NfL, using the highly sensitive Simoa assay, in samples from the UK SBMA cohort, and compared them with healthy controls and with slow and fast progressor ALS cases, as positive controls. While fast and slow ALS progressors showed the previously described¹¹ increase in NfL levels ($p < 0.0001$ and $p = 0.0008$, respectively), we detected no significant increase in the SBMA cases ($p = 0.99$). This finding was replicated in the Italian SBMA cohort

Table Cohort demographic and genetic information, neurofilament light chain (NfL), creatine kinase (CK), and creatinine levels

Patient group	n	Age, y, mean (SD)	Sex, F/M	AMAT	SBMAFRS	ALSFRS, mean (SD)	Disease duration onset–last visit, mo, mean (SEM)	CAG repeats	NfL, pg/mL (SEM)	CK, U/L (SEM)	Creatinine, μ mol/L (SEM)
HC UK	74	60.04 (8.7)	31/43	—	—	—	—	—	29 (2.2), n = 50	273 (31), n = 43 all male	88 (9.5), n = 6 all male
SBMA UK	50	56.35 (11.1)	0/50	31 (1.48), n = 43	41 (1.27), n = 42	38 (0.80), n = 43	—	45.5 (40–53)	19 (1.59), n = 48	951 (88.0), n = 46	54 (2.62), n = 46
HC Italy	23	NA	NA	—	—	—	—	—	28 (5.8)	NA	NA
SBMA Italy	43	59 (10)	0/43	24 (1.91), n = 32	42 (1.12), n = 35	NA	—	45.8 (41–51)	25 (1.16)	940 (107)	52 (2.14), n = 41
Fast ALS UK	40	65 (8)	16/24	—	—	31.8 (8.6), n = 39	11 (1.2), n = 26	NA	365 (41.0), n = 27	307 (73.6), n = 17 all male	76 (5.1), n = 4 all male
Slow ALS UK	62	63 (14)	8/54	—	—	39.6 (7.3), n = 60	62 (11.4), n = 26	NA	112 (24.7), n = 26	274 (31.2), n = 43 all male	82 (5.8), n = 11 all male

Abbreviations: ALS = amyotrophic lateral sclerosis; ALSFRS = Amyotrophic Lateral Sclerosis Functional Rating Scale; AMAT = Adult Myopathy Assessment Tool; HC = healthy controls; SBMA = spinal and bulbar muscular atrophy; SBMAFRS = Spinal and Bulbar Muscular Atrophy Functional Rating Scale.

($p > 0.99$) (figure 1A). We further examined whether levels changed longitudinally at 12 ($n = 28$) and 24 ($n = 8$) months, and showed NfL levels to be stable across this time, with the exception of 3/28 cases where levels increased to >58 pg/mL, the 95th percentile of controls (figure 1B).

NfL levels are unchanged in a mouse model of disease

We assessed NfL levels in a well-established SBMA mouse model: AR100 mice. These mice express a 100-CAG expansion in the human AR and develop a late-onset progressive neuromuscular phenotype characterized by muscle atrophy and motor neuron degeneration, with no motor neuron loss at 6 months, and a 40% loss of motor neurons at 18 months of age (B. Malik and L. Greensmith, personal written communication, July 30, 2018, and references 12 and 17). NfL measurements performed on the same platform as for the human samples did not show a significant increase in AR100 mice at 18 months vs littermate controls ($n = 9$; $p = 0.45$) (figure 1C).

NfL levels are not increased in blood-derived biofluids from 2 independent SBMA cohorts and a well-established mouse model of disease.

Muscle damage and muscle mass markers are significantly altered in SBMA

We investigated the levels of markers of muscle damage (CK) and muscle mass (creatinine) in our patient cohorts using standard clinical diagnostic assays. CK levels showed remarkably similar values in both the UK and Italian cohorts (mean 951 and 940 U/L, respectively), and were significantly increased compared to both healthy controls (mean 122 U/L, $p < 0.0001$ for both UK and Italian cohorts) and ALS ($p < 0.0001$ for both UK and Italian cohorts vs slow and fast progressing ALS). Interestingly, CK levels, differently from NfL, did not differ between slow and fast progressing ALS (mean 699 and 701 U/L, respectively, $p > 0.99$) (figure 2A).

Creatinine levels were very similar in UK and Italian cohorts and significantly decreased compared to controls and ALS (figure 2C).

Creatinine levels correlate with SBMA severity

To evaluate disease severity, we used the SBMAFRS, a function rating scale developed for SBMA, and the AMAT score, which detects deficits in functional muscle performance and endurance and has been used as an outcome measure for SBMA clinical trials.^{13–15} While no significant correlation was present between clinical measures and CK (figure 2B), creatinine levels strongly inversely correlated with both SBMAFRS and AMAT in both SBMA cohorts (figure 2D).

Discussion

We show that NfL concentration in blood is unchanged in patients with SBMA compared with controls, and concentrations do not change over time and are not related to disease severity. The result was confirmed in a well-established mouse model of SBMA. This is surprising given that NfL has been described to be increased in numerous neurodegenerative conditions and mouse models, and cannot simply be imputed solely to the slowly progressive nature of SBMA, as a recent study using the same technique has detected an NfL increase in Charcot-Marie-Tooth disease, another very slowly progressive neuromuscular condition.^{10,11,16,18,19}

Conversely, markers of muscle damage (CK) and muscle mass (creatinine) are aberrant in SBMA. CK is elevated compared to both controls and ALS cases, and does not correlate to clinical measures, possibly due to its reflection of more acute muscle damage, whereas creatinine is reduced and significantly correlates with disease severity. A decrease in creatinine has also been shown in a Japanese SBMA cohort to correlate with grip strength and 6-minute walk distance and has also been shown

Figure 1 Neurofilament light chain (NfL) levels are unchanged in spinal and bulbar muscular atrophy (SBMA)

(A) NfL concentrations (pg/mL) from the UK cohort (plasma, UK control [CTR], UK SBMA, slow amyotrophic lateral sclerosis [ALS], and fast ALS) and the Italian cohort (IT CTR and IT SBMA). Assays were conducted together, but statistical analysis (analysis of variance, Bonferroni multiple comparison correction) is represented only between samples from the same cohort. (B) NfL concentrations in the longitudinal study in the UK cohort. Twelve-month ($n = 28$) and 24-month ($n = 8$) timepoints represent 12 ± 2 and 24 ± 2 months. (C) NfL levels from AR100 (SBMA) and littermate control (WT) mice are shown (Mann-Whitney test). Columns indicate mean, error bars indicate SD; *** $p < 0.001$; **** $p < 0.0001$; ns $p > 0.05$.

Figure 2 Creatine kinase (CK) and creatinine levels are altered in spinal and bulbar muscular atrophy (SBMA) and correlate with clinical severity

(A) CK SBMA plasma concentrations from both UK and Italian cohorts are significantly increased compared to controls (CTR) ($p < 0.0001$) and slow ($p < 0.0001$) and fast ($p = 0.0003$, United Kingdom; $p = 0.0002$, Italy) progressive amyotrophic lateral sclerosis (ALS). (B) CK concentrations of both UK and Italian cohorts do not significantly correlate with the Spinal and Bulbar Muscular Atrophy Functional Rating Scale (SBMAFRS) (left) and Adult Myopathy Assessment Tool (AMAT) (right) clinical severity scores (for SBMAFRS: United Kingdom, $r = -0.02$, $p = 0.99$; Italian, $r = 0.25$, $p = 0.15$; and for AMAT: United Kingdom, $r = 0.15$, $p = 0.32$; IT, $r = 0.29$, $p = 0.11$). (C) Creatinine plasma concentrations from both UK and Italian cohorts are significantly decreased compared to controls ($p < 0.0001$) and slow progressive ALS ($p < 0.0001$). (D) Creatinine concentrations of both UK and Italian cohorts significantly correlate with the SBMAFRS (left) and AMAT (right) clinical severity scores (for SBMAFRS: United Kingdom, $r = 0.61$, $p < 0.0001$; Italy, $r = 0.49$, $p = 0.0035$; and for AMAT: United Kingdom, $r = 0.75$, $p < 0.0001$; Italy, $r = 0.50$, $p = 0.0044$). Blue (United Kingdom) and black (Italy) lines represent the best-fit line. Red dashed lines represent upper and lower extremities of the normal range. Columns indicate mean, error bars indicate SD; **** $p < 0.0001$; ns $p > 0.05$.

to be lower in patients with SBMA than in patients with ALS who have comparable wasting of muscle mass.^{20,21}

Traditionally SBMA has been considered a lower motor neuron (LMN) disease. Indeed, denervation is a crucial neurophysiology finding, and loss of LMN in postmortem tissue has been well-documented.^{2,22} Muscle abnormalities and CK levels elevated beyond those found in ALS have been described in early case series of this disease.²³ Nonetheless, only recently, analysis of SBMA muscle biopsies, and functional work specifically exploring the role of muscle in mouse models, have led to acknowledging a primary muscle component to be relevant for disease progression and pathogenesis alongside the neurogenic component,^{3–9} and our findings support this.

Finally, SBMA and ALS are in differential diagnosis, and a substantial proportion of patients with SBMA initially receive a diagnosis of ALS, causing distress.²⁴ Although genetic clinical testing is the gold standard for SBMA diagnosis, the finding of normal levels of NfL in SBMA could be used in a diagnostic panel of biochemical markers to help differentiate patients presenting with motor neuron deficits.

NfL concentrations in blood were unchanged in 2 cohorts of patients with SBMA and in a mouse model of disease, while markers of muscle damage and mass were altered, the latter showing correlation with clinical measures of disease, suggesting that biomarkers of muscle damage and mass, rather than neuronal damage, should be used to monitor disease progression and outcome. This is consistent with previous observations that a primary myopathic component plays a primary role in the disorder^{3–9} and supports the development of novel disease-modifying agents towards the muscle target and the incorporation of muscle biomarkers to objectively assess outcomes in therapeutic trials in SBMA.

Acknowledgment

The authors thank the patients involved in the study and their families for participation in and support of KD research.

Study funding

P.F. is supported by an MRC/MNDA LEW Clinician Scientist Fellowship (MR/M008606/1) and the NIHR UCLH Biomedical Research Centre. This study was supported by the NIHR UCLH Biomedical Research Centre, the UCL

Kennedy's Disease Fund, and KDUK. O.J.Z. is funded by the UK National Institute of Health Research on an Academic Clinical Fellowship. G.S. is supported by Grant Telethon GTB12001D (Telethon Biobank ed EuroBiobank), Telethon GUP15009B, and AFM-Telethon grant number 18722. C.R. is funded by a Wellcome Trust Clinical Research Career Development Fellowship and the Muscular Dystrophy Association (MDA). H.Z. is funded by the UK Dementia Research Institute at UCL, the European Research Council, the Swedish Research Council, and the Knut and Alice Wallenberg Foundation. The Simoa instrument was bought using a Wellcome Trust multi-user equipment grant (PI: H.Z.). Study Funded by the NIHR UCLH Biomedical Research Centre Grant #BRC279566.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication history

Received by *Neurology* August 12, 2018. Accepted in final form November 5, 2018.

Appendix Authors

Name	Location	Role	Contribution
Vittoria Lombardi, PhD	Blizard Institute, Queen Mary, University of London, UK	Author	Data acquisition and data analysis, statistical analysis, writing of the manuscript
Giorgia Querin, MD	Department of Neurosciences, University of Padova, Italy	Author	Data acquisition and data analysis
Luca Zampedri	Institute of Neurology, University College London, UK	Author	Data acquisition and data analysis
Oliver J. Ziff, MD, MRCP	Institute of Neurology, University College London, UK	Author	Statistical analysis, writing of the manuscript
Caroline Heller	Institute of Neurology, University College London, UK	Author	Data acquisition and data analysis
Martha Foiani	Institute of Neurology, University College London, UK	Author	Data acquisition and data analysis
Ilaria Martinelli, MD	Department of Neurosciences, University of Padova, Italy	Author	Data acquisition and data analysis
Cinzia Bertolin, PhD	Department of Neurosciences, University of Padova, Italy	Author	Data acquisition and data analysis
Ching-Hua Lu, MD, PhD	Department of Neurology, China Medical University Hospital, Taichung City, Taiwan	Author	Data acquisition and data analysis

Appendix (continued)

Name	Location	Role	Contribution
Bilal Malik, PhD	Institute of Neurology, University College London, UK	Author	Data acquisition and data analysis
Kezia Allen	Basildon Hospital, UK	Author	Data acquisition and data analysis
Carlo Rinaldi, MD, PhD	University of Oxford, UK	Author	Data acquisition and data analysis, critical revision of the manuscript
Linda Greensmith, PhD	Institute of Neurology, University College London, UK	Author	Critical revision of the manuscript
Henrik Zetterberg, PhD	Institute of Neurology, University College London, UK	Author	Critical revision of the manuscript
Amanda Heslegrave, PhD	Institute of Neurology, University College London, UK	Author	Data acquisition and data analysis, critical revision of the manuscript
Linda Greensmith, PhD	Institute of Neurology, University College London, UK	Author	Critical revision of the manuscript
Michael Hanna, FRCP	Institute of Neurology, University College London, UK	Author	Critical revision of the manuscript
Gianni Soraru, MD, PhD	Department of Neurosciences, University of Padova, Italy	Author	Data acquisition and data analysis, critical revision of the manuscript
Andrea Malaspina, MD, PhD	Blizard Institute, Queen Mary, University of London, UK	Author	Designed and conceptualized study, data acquisition and data analysis, critical revision of the manuscript
Pietro Fratta, MD, PhD	Institute of Neurology, University College London, UK	Author	Designed and conceptualized study, data acquisition and data analysis, statistical analysis, writing of the manuscript

References

- Cortes CJ, La Spada AR. X-linked spinal and bulbar muscular atrophy: from clinical genetic features and molecular pathology to mechanisms underlying disease toxicity. *Adv Exp Med Biol* 2018;1049:103–133.
- Kennedy WR, Alter M, Sung JH. Progressive proximal spinal and bulbar muscular atrophy of late onset: a sex-linked recessive trait. *Neurology* 1968;18:671–680.
- Cortes CJ, Ling SC, Guo LT, et al. Muscle expression of mutant androgen receptor accounts for systemic and motor neuron disease phenotypes in spinal and bulbar muscular atrophy. *Neuron* 2014;82:295–307.
- Giorgetti E, Yu Z, Chua JP, et al. Rescue of metabolic alterations in AR113Q skeletal muscle by peripheral androgen receptor gene silencing. *Cell Rep* 2016;17:125–136.
- Johansen JA, Yu Z, Mo K, et al. Recovery of function in a myogenic mouse model of spinal bulbar muscular atrophy. *Neurobiol Dis* 2009;34:113–120.
- Soraru G, D'Ascenzo C, Polo A, et al. Spinal and bulbar muscular atrophy: skeletal muscle pathology in male patients and heterozygous females. *J Neurol Sci* 2008;264:100–105.
- Malena A, Pennuto M, Tezze C, et al. Androgen-dependent impairment of myogenesis in spinal and bulbar muscular atrophy. *Acta Neuropathol* 2013;126:109–121.
- Monks DA, Johansen JA, Mo K, et al. Overexpression of wild-type androgen receptor in muscle recapitulates polyglutamine disease. *Proc Natl Acad Sci USA* 2007;104:18259–18264.
- Chahin N, Sorenson EJ. Serum creatine kinase levels in spinobulbar muscular atrophy and amyotrophic lateral sclerosis. *Muscle Nerve* 2009;40:126–129.

10. Sandelius Å, Zetterberg H, Blennow K, et al. Plasma neurofilament light chain concentration in the inherited peripheral neuropathies. *Neurology* 2018;90:e518–e524.
11. Lu C-H, Macdonald-Wallis C, Gray E, et al. Neurofilament light chain: a prognostic biomarker in amyotrophic lateral sclerosis. *Neurology* 2015;84:2247–2257.
12. Sopher BL, Thomas PS, LaFevre-Bernt MA, et al. Androgen receptor YAC transgenic mice recapitulate SBMA motor neuronopathy and implicate VEGF164 in the motor neuron degeneration. *Neuron* 2004;41:687–699.
13. Harris-Love MO, Fernandez-Rhodes L, Joe G, et al. Assessing function and endurance in adults with spinal and bulbar muscular atrophy: validity of the adult myopathy assessment tool. *Rehabil Res Pract* 2014;2014:873872.
14. Querin G, DaRe E, Martinelli I, et al. Validation of the Italian version of the SBMA Functional Rating Scale as outcome measure. *Neurol Sci* 2016;37:1815–1821.
15. Hashizume A, Katsuno M, Suzuki K, et al. A functional scale for spinal and bulbar muscular atrophy: cross-sectional and longitudinal study. *Neuromuscul Disord* 2015;25:554–562.
16. Lu CH, Petzold A, Kalmar B, Dick J, Malaspina A, Greensmith L. Plasma neurofilament heavy chain levels correlate to markers of late stage disease progression and treatment response in SOD1(G93A) mice that model ALS. *PLoS One* 2012;7:e40998.
17. Malik B, Nirmalanathan N, Bilsland LG, et al. Absence of disturbed axonal transport in spinal and bulbar muscular atrophy. *Hum Mol Genet* 2011;20:1776–1786.
18. Rohrer JD, Woollacott IOC, Dick KM, et al. Serum neurofilament light chain protein is a measure of disease intensity in frontotemporal dementia. *Neurology* 2016;87:1329–1336.
19. Bacioglu M, Maia LF, Preische O, et al. Neurofilament light chain in blood and CSF as marker of disease progression in mouse models and in neurodegenerative diseases. *Neuron* 2016;91:56–66.
20. Hijikata Y, Katsuno M, Suzuki K, et al. Impaired muscle uptake of creatine in spinal and bulbar muscular atrophy. *Ann Clin Transl Neurol* 2016;3:537–546.
21. Hashizume A, Katsuno M, Banno H, et al. Longitudinal changes of outcome measures in spinal and bulbar muscular atrophy. *Brain* 2012;135:2838–2848.
22. Ogata A, Matsuura T, Tashiro K, et al. Expression of androgen receptor in X-linked spinal and bulbar muscular atrophy and amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 1994;57:1274–1275.
23. Harding AE, Thomas PK, Baraitser M, Bradbury PG, Morgan-Hughes JA, Ponsford JR. X-linked recessive bulbospinal neuronopathy: a report of ten cases. *J Neurol Neurosurg Psychiatry* 1982;45:1012–1019.
24. Fratta P, Nirmalanathan N, Masset L, et al. Correlation of clinical and molecular features in spinal bulbar muscular atrophy. *Neurology* 2014;82:2077–2084.

Neurology®

Muscle and not neuronal biomarkers correlate with severity in spinal and bulbar muscular atrophy

Vittoria Lombardi, Giorgia Querin, Oliver J. Ziff, et al.

Neurology 2019;92:e1205-e1211 Published Online before print February 20, 2019

DOI 10.1212/WNL.0000000000007097

This information is current as of February 20, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/92/11/e1205.full
References	This article cites 24 articles, 8 of which you can access for free at: http://n.neurology.org/content/92/11/e1205.full#ref-list-1
Citations	This article has been cited by 4 HighWire-hosted articles: http://n.neurology.org/content/92/11/e1205.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Anterior nerve cell disease http://n.neurology.org/cgi/collection/anterior_nerve_cell_disease
Errata	An erratum has been published regarding this article. Please see next page or: /content/94/19/852.1.full.pdf
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

Disputes & Debates: Editors' Choice

Steven Galetta, MD, FAAN, Section Editor

Editors' note: Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source

In "Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source," Ntaios et al. studied 777 patients with embolic stroke of undetermined source (ESUS) and found that the presence of ipsilateral nonstenotic carotid plaque was associated with a significantly lower likelihood for atrial fibrillation. Siegler et al. questioned the modality used to identify nonstenotic carotid plaque and paroxysmal atrial fibrillation, given that this affects incidence rates. Furthermore, they noted that although the authors commented on the frequency of conditions during which atrial fibrillation was detected (such as routine ECG, stroke recurrence, and myocardial infarction), it would have been useful to compare the incidence rates for patients with and without plaque. Ntaios responded that unfortunately, they did not have access to information about the technique used to diagnose carotid plaque or atrial fibrillation and that they did not think it was appropriate to compare the incidence of conditions during which atrial fibrillation was detected without this information. Lattanzi et al. commented that it would be useful to assess the relationship between plaques and atrial fibrillation and other variables, such as atrial cardiopathy, which has been shown to be inversely related to artery-to-artery strokes. Ntaios responded that they have performed additional analysis and hope to have their results published soon. It is clear that further research into workup and management of ESUS is needed.

Ariane Lewis, MD, and Steven Galetta, MD
Neurology® 2020;94:849. doi:10.1212/WNL.0000000000009426

Reader response: Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source

James E. Siegler (Philadelphia), Jesse Thon (Philadelphia), and Brett L. Cucchiara (Philadelphia)
Neurology® 2020;94:849–850. doi:10.1212/WNL.0000000000009428

We read with much interest the multicenter observational experience of Ntaios et al.¹ regarding the incidence of atrial fibrillation among patients with embolic stroke of undetermined source (ESUS). The article added considerably to the increasing evidence regarding the heterogeneity of ESUS and the clinical relevance of nonstenotic carotid artery plaque.^{2,3} It may also assist in decision making regarding outpatient cardiac event monitoring.

We were curious to know whether the investigators could expand on their variable definitions. For the primary independent variable of nonstenotic ipsilateral carotid plaque, the various imaging modalities used could lead to inaccurate prevalence estimates. It would be helpful to know which modalities were used in patients with plaque vs those without. Furthermore, were the methods used for detecting paroxysmal atrial fibrillation (pAF) similar between these groups? At our center, we routinely monitor for pAF for a minimum of 14 days in patients with ESUS older than 50 years—irrespective of the presence of nonstenotic plaque. But we confess to our own selection bias that longer monitoring may be pursued in patients who have minimal or no plaque and in older patients. As has been described previously,⁴ the device and duration of

event monitoring can dramatically affect the probability of detecting pAF in patients with cryptogenic stroke. The investigators presented data on conditions for which pAF was identified (table 2),¹ but it would be more important to compare the incidence rates for these conditions between patients with plaque and those without.

1. Ntaios G, Perlepe K, Sirimarco G, et al. Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source. *Neurology* 2019;92:e2644–e2652.
2. Freilinger TM, Schindler A, Schmidt C, et al. Prevalence of nonstenosing, complicated atherosclerotic plaques in cryptogenic stroke. *JACC Cardiovasc Imaging* 2012;5:397–405.
3. Coutinho JM, Derkatch S, Potvin AR, et al. Nonstenotic carotid plaque on CT angiography in patients with cryptogenic stroke. *Neurology* 2016;87:665–672.
4. Choe WC, Passman RS, Brachmann J, et al. A comparison of atrial fibrillation monitoring strategies after cryptogenic stroke (from the cryptogenic stroke and underlying AF trial). *Am J Cardiol* 2015;116:889–893.

Author response: Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source

George Ntaios (Larissa, Greece)

Neurology® 2020;94:850. doi:10.1212/WNL.0000000000009427

I would like to thank Dr. Siegler and colleagues for their kind words and for their thoughtful comments on our article.¹ I agree that it would be useful to know which imaging modalities were used to assess carotid plaques; however, we did not register this information. Furthermore, we did not assess the type of modalities of heart rhythm monitoring that applied to our patients and therefore cannot provide information on whether they were used similarly between patients with or without carotid stenosis. I agree that it may be possible that these modalities were applied differently between these 2 groups. In this context, any comparison between these 2 groups about the incidence rates of the conditions during which atrial fibrillation was detected could lead to erroneous conclusions, which is why we refrained from providing these results in the article.¹

1. Ntaios G, Perlepe K, Sirimarco G, et al. Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source. *Neurology* 2019;92:e2644–e2652.

Copyright © 2020 American Academy of Neurology

Reader response: Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source

Simona Lattanzi (Ancona, Italy) and Mauro Silvestrini (Ancona, Italy)

Neurology® 2020;94:850–851. doi:10.1212/WNL.0000000000009430

We read with great interest the article by Ntaios et al.,¹ which demonstrated a lower cumulative probability of atrial fibrillation (AF) detection in patients with embolic strokes of undetermined source (ESUS) with ipsilateral nonstenotic carotid plaques compared with those without.

Far from being a single entity, ESUS constitute a heterogeneous group with high variability in clinical presentation and etiology. Remarkably, left atrialopathy has been inversely associated with paradoxical and artery-to-artery embolic sources, including patent foramen ovale (PFO) and vulnerable substenotic atherosclerotic disease of aortic arch and neck arteries.² Carotid plaques with increased thickness, mobility, and ulceration were more common when ipsilateral than contralateral to the stroke site.³ Young patients presented lower rates of vascular risk factors and left atrial enlargement, higher incidence of PFO, and no atherosclerosis.⁴

Accordingly, it would be interesting to provide a more comprehensive analysis of data by exploring the relationships between nonstenotic carotid plaques and AF detection with additional parameters, including demographics, stroke severity and location, indicators of plaque vulnerability, presence of atrial cardiopathy, and paradoxical embolic sources. Optimal management for ESUS is to be elucidated. Efforts to identify distinct phenotypes based on the underlying pathogenesis could inform tailored strategies and improve prevention.⁵

1. Ntaios G, Perlepe K, Sirimarco G, et al. Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source. *Neurology* 2019;92:e2644–e2652.
2. Lattanzi S, Cagnetti C, Pulcini A, et al. The P-wave terminal force in embolic strokes of undetermined source. *J Neurol Sci* 2017;375:175–178.
3. Komatsu T, Iguchi Y, Arai A, et al. Large but nonstenotic carotid artery plaque in patients with a history of embolic stroke of undetermined source. *Stroke* 2018;49:3054–3056.
4. Piffer S, Bignamini V, Rozzanigo U, et al. Different clinical phenotypes of embolic stroke of undetermined source: a subgroup analysis of patients. *J Stroke Cerebrovasc Dis* 2018;27:3578–3586.
5. Lattanzi S, Brigo F, Cagnetti C, et al. Patent foramen ovale and cryptogenic stroke or transient ischemic attack: to close or not to close? A systematic review and meta-analysis. *Cerebrovasc Dis* 2018;45:193–203.

Copyright © 2020 American Academy of Neurology

Author response: Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source

George Ntaios (Larissa, Greece)

Neurology® 2020;94:851. doi:10.1212/WNL.00000000000009429

I thank Dr. Lattanzi and Dr. Silvestrini for their interest in our work¹ and their constructive suggestions. Several of the proposed analyses were already conducted and are currently under review in peer-reviewed journals. Hopefully, the results of these analyses will be publicly available soon.

1. Ntaios G, Perlepe K, Sirimarco G, et al. Carotid plaques and detection of atrial fibrillation in embolic stroke of undetermined source. *Neurology* 2019;92:e2644–e2652.

Copyright © 2020 American Academy of Neurology

Disputes & Debates: Rapid online correspondence

The editors encourage comments on recent articles through Disputes & Debates:

Access an article at *Neurology.org/N* and click on “MAKE COMMENT” beneath the article header. Responses will be posted as rapidly as possible.

Before submitting a comment to Disputes & Debates, remember the following:

- Disputes & Debates is restricted to comments about articles published in *Neurology* within the last 8 weeks
 - Read previously posted comments; redundant comments will not be posted
 - Your submission must be 200 words or less and have a maximum of 5 references; the first reference must be the article on which you are commenting
 - You can include a maximum of 5 authors (including yourself)
-

Author disclosures are available upon request (journal@neurology.org).

Muscle and not neuronal biomarkers correlate with severity in spinal and bulbar muscular atrophy

Neurology® 2020;94:852. doi:10.1212/WNL.0000000000009392

In the article “Muscle and not neuronal biomarkers correlate with severity in spinal and bulbar muscular atrophy” by Lombardi et al.,¹ the y-axis label for figure 2c should be “Creatinine (umol/L).” The editorial office regrets the error.

Reference

1. Lombardi V, Querin G, Ziff OJ, et al. Muscle and not neuronal biomarkers correlate with severity in spinal and bulbar muscular atrophy. *Neurology* 2019;92:e1205–e1211.

Finding a common path to the assessment of persons with intellectual development disorders

Neurology® 2020;94:852. doi:10.1212/WNL.0000000000009541

In the editorial “Finding a common path to the assessment of persons with intellectual development disorders” by Coffman and Borgatti,¹ Dr. Borgatti’s affiliation should read: “Child Neurology and Psychiatry Unit, IRCCS Mondino Foundation (Pavia, Italy).” The authors regret the error.

Reference

1. Coffman KA, Borgatti R. Finding a common path to the assessment of persons with intellectual development disorders. *Neurology* 2020;94:507–508.

Visual field deficits following laser ablation of the hippocampus

Neurology® 2020;94:852. doi:10.1212/WNL.0000000000009543

In the article “Visual field deficits following laser ablation of the hippocampus” by Donos et al.,¹ first published online February 26, 2020, the figures should appear in the correct numbered order and should correctly match the numbered figure legends. The image under figure 1 should have appeared with figure 3. The image under figure 2 should have appeared under figure 4. The image for figure 3 should have corresponded with figure 5. The image under figure 4 should have appeared with figure 6. The image under figure 5 should have appeared under figure 1. The image under figure 6 should have appeared under figure 2. The figures and legends are correctly matched in the March 24, 2020, issue. The publisher regrets the error.

Reference

1. Donos C, Rollo P, Tombridge K, Johnson JA, Tandon N. Visual field deficits following laser ablation of the hippocampus. *Neurology* 2020;94:e1303–e1313.