Visibly enlarged peripheral nerves in lepromatous leprosy

Joy Zhuo Ding, MD, and Pedro Legua, MD

Neurology® 2019;92:627-628. doi:10.1212/WNL.00000000000007239

Correspondence

Dr. Ding
joy.ding@medportal.ca

A 52-year-old man from the Peruvian jungle presented with a 4-year history of progressive distal numbness and facial swelling. He had diffusely infiltrated skin on his face, torso, and extremities (figure, A and B), with patchy pinprick sensation loss. Multiple peripheral nerves were palpable, including ulnar and fibular nerves, in addition to visibly enlarged great auricular (figure, C) and superficial radial nerves (figure, D).

Slit skin smears showed up to 5+ acid-fast bacilli. Skin biopsy was compatible with lepromatous leprosy, strongly positive with Fite-Faraco staining.

Enlargement of peripheral nerves in leprosy is well-described; however, hypertrophy visible to the naked eye is rarely photographed.

Author contributions

J.Z. Ding: manuscript conception, drafting, and editing and taking clinical photographs.
P. Legua: manuscript editing and clinical care of the patient.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

From the Division of Neurology (J.Z.D.), The Ottawa Hospital, Canada; Instituto de Medicina Tropical "Alexander von Humboldt" (P.L.), Universidad Peruana Cayetano Heredia; and Departamento de Enfermedades Infecciosas, Tropicales y Dermatológicas (P.L.), Hospital Cayetano Heredia, Lima, Peru.

Go to Neurology.org/N for full disclosures.

(A) Diffusely infiltrated and erythematous skin. (B) Diffuse infiltration of the face with loss of eyebrows. (C) Visibly enlarged greater auricular nerve; also note the enlarged earlobe. (D) Visibly enlarged superficial radial nerve.

Are You Ready?

The MACRA's new Quality Payment Program is now in effect. Learn about your payment and reporting options under the Merit-based Incentive Payment System (MIPS) and Advanced Alternative Payment Models (APMs) and what you need to know to do now to ensure your success. Learn more at [AAN.com/view/QPP](https://www.aan.com/view/QPP) and sign up to receive timely updates.

Neurology®

Visibly enlarged peripheral nerves in lepromatous leprosy

Joy Zhuo Ding and Pedro Legua

Neurology 2019;92;627-628

DOI 10.1212/WNL.00000000000007239

This information is current as of March 25, 2019

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/92/13/627.full>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All global neurology

http://n.neurology.org/cgi/collection/all_global_neurology

Bacterial infections

http://n.neurology.org/cgi/collection/bacterial_infections

Clinical neurology examination

http://n.neurology.org/cgi/collection/clinical_neurology_examination

EMG

<http://n.neurology.org/cgi/collection/emg>

Peripheral neuropathy

http://n.neurology.org/cgi/collection/peripheral_neuropathy

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.