Teaching NeuroImages: Greater occipital nerve injection

A cautionary tale

Diana Y. Wei, MBBS, MRCP, Steve Connor, MRCP, FRCR, and Peter J. Goadsby, MD, PhD

Neurology® 2019;92:e746-e747. doi:10.1212/WNL.0000000000006929

Correspondence

Dr. Goadsby
peter.goadsby@kcl.ac.uk

Figure 1 Lateral and frontal skull radiographs

(A) Lateral. (B) Frontal. The occipital craniectomy is demonstrated on the lateral projection (arrow) but partially obscured by skull base and partially excluded from the image. Radiographs are less expensive and involve lower radiation dose compared with CT imaging; however, skull base surgical defects may not be optimally demonstrated.

A 29-year-old man with medically refractory right-sided chronic cluster headache presented with increased attack frequency. As a bridging therapy, we considered a greater occipital nerve (GON) injection with corticosteroids and local anesthetic. Delving through his childhood medical records, it came to light he had an incidental right cerebellopontine angle dermoid excised. He could not recall the details of this; therefore, plain skull X-ray (figure 1) and head CT (figure 2) were performed. Neuroimaging is required before proceeding with GON injection if there is any suggestion of past neurosurgical intervention. There is a risk of causing loss of consciousness with GON injections into nonintact skull.^{1,2}

MORE ONLINE

→Teaching slides

links.lww.com/WNL/A813

Author contributions

D.Y. Wei: case report concept, wrote the initial manuscript, acquisition of data, review of manuscript. S. Connor: interpretation of X-ray and CT, created CT reconstruction, review of manuscript. P.J. Goadsby: case report concept, critical revision of manuscript for intellectual content.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

- Okuda Y, Matsumoto T, Shinohara M, Kitajima T, Kim P. Sudden unconsciousness during a lesser occipital nerve block in a patient with the occipital bone defect. *Eur J Anaesthesiol* 2001;18:829–832.
- Sprenger T, Seifert CL. Coma after greater occipital nerve blockade in a patient with previous posterior fossa surgery. *Headache* 2013;53:548–550.

From the Headache Group, Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience (D.Y.W., P.J.G.), and School of Biomedical Engineering and Imaging Sciences (S.C.), King's College London; and Departments of Neuroradiology (S.C.) and Neurology (P.J.G.), King's College Hospital NHS Foundation Trust, London, UK.

Figure 2 Oblique posterior view of a volume-rendered reconstruction of the CT data

The defect within the right occipital bone at the site of the craniectomy is demonstrated.

Neurology®

Teaching NeuroImages: Greater occipital nerve injection: A cautionary tale

Diana Y. Wei, Steve Connor and Peter J. Goadsby

Neurology 2019;92:e746-e747

DOI 10.1212/WNL.0000000000006929

This information is current as of February 11, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/92/7/e746.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/92/7/e746.full#ref-list-1
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.