Teaching NeuroImages: Sleep-onset REM period during routine EEG

Karl A. Kasischke, MD, Amanda Pennington, MD, PhD, and Selim R. Benbadis, MD

Neurology® 2019;93:e1123-e1124. doi:10.1212/WNL.0000000000008108

Correspondence

Dr. Kasischke
kkasischke@health.usf.edu

MORE ONLINE

→Teaching slides

links.lww.com/WNL/A950

Figure Two consecutive EEG pages in bilateral longitudinal (double banana) montage

(A) EEG sample with sleep spindles in normal stage N2 sleep. (B) On the contiguous second page, there is a burst of high-amplitude lateral eye movements with concurrent sawtooth waves, indicative of REM. Also note muscle quiescence. This sleep-onset REM period could be misinterpreted as sharp waves.

A 76-year-old incarcerated man presented with word-finding difficulties and underwent a routine EEG. Shortly after reaching stage N2 sleep, EEG showed brisk lateral eye movements with concurrent sawtooth waves (figure), indicating sleep-onset REM period (SOREMP). Normal REM sleep is reached 90–120 minutes after sleep onset. SOREMP is an abnormal EEG

From the Department of Neurology, Morsani College of Medicine, University of South Florida, Tampa.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

finding thought to be suggestive of narcolepsy. However, it may also be present in 3.9% of the general population¹ and is normal in infants. It is also associated with severe sleep deprivation, obstructive sleep apnea, and alcohol or medication (selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, tricyclic antidepressants, monoamine oxidase inhibitors) withdrawal. In the inpatient setting, it is usually caused by sleep deprivation.²

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Appendix Authors

Name	Location	Role	Contribution
Karl A. Kasischke, MD	University of South Florida, Tampa	Author	Analyzed data, drafted and revised the manuscript
Amanda Pennington, MD, PhD	University of South Florida, Tampa	Author	Analyzed and presented data
Selim R. Benbadis, MD	University of South Florida, Tampa	Author	Designed study and revised the manuscript

References

1. Sing M, Drake CL, Roth T. The prevalence of multiple sleep-onset REM periods in a population-based sample. *Sleep* 2006;29:890–895.

2. Gangadhara S, Pizarro-Otero J, Bozorg A, Benbadis S. The significance of REM sleep on routine EEG. *Neurodiagn J* 2016;56:37–40.

Neurology®

Teaching NeuroImages: Sleep-onset REM period during routine EEG

Karl A. Kasischke, Amanda Pennington and Selim R. Benbadis

Neurology 2019;93:e1123-e1124

DOI 10.1212/WNL.00000000000008108

This information is current as of September 9, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/93/11/e1123.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/93/11/e1123.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All clinical neurophysiology http://n.neurology.org/cgi/collection/all_clinical_neurophysiology EEG http://n.neurology.org/cgi/collection/eeg Narcolepsy http://n.neurology.org/cgi/collection/narcolepsy
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.