Teaching NeuroImages: Frontal lobe involvement in adult-onset cerebral X-linked adrenoleukodystrophy

Chenchen Liu, MD,* Yayun Cao, MD,* and Bitao Bu, MD

Neurology® 2019;93:e1326-e1327. doi:10.1212/WNL.00000000000008181

Correspondence

Dr. Bu
bubitao@yeah.net

MORE ONLINE

→Teaching slides

links.lww.com/WNL/A960

A 42-year-old man presented with 1 year of progressive behavioral and personality changes. His antenatal, birth, and neurodevelopment were unremarkable. Examination showed hyperactive behavior, lower limbs paralysis (Medical Research Council grading 3) with pyramidal signs, and no skin hyperpigmentation or visual disturbance. Brain MRI revealed diffuse lesions in both frontal lobes, with peripheral rim restricted diffusion and enhancement (figure 1). Elevated plasma very-long-chain fatty acid level, p.Ser98Ter mutation in *ABCD1* gene, and his pedigree (figure 2) confirmed

Figure 1 Brain MRI

Axial T2-weighted (A) and fluid-attenuated inversion recovery (B) images show diffuse white matter hyperintensity in both frontal lobes. Diffusion-weighted imaging sequence reveals peripheral rim restricted diffusion in the white matter of the frontal lobes (C), with enhancement (D).

*These authors contributed equally to this work.

From the Department of Neurology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Figure 2 Genetic testing and family tree

(A) Sequence determination of c.293C>A (p.Ser98Ter) mutation of the *ABCD1* gene. The arrow indicates the involved nucleotide. (B) Family tree: The presented case is marked with an arrow. III-2, III-4, and IV-8 had X-linked adrenoleukodystrophy; I-2, II-1, II-4, III-8, IV-2, IV-3, and IV-9 were female carriers.

the diagnosis of adult-onset cerebral X-linked adrenoleukodystrophy (X-ALD). X-ALD is an inborn error of metabolism predominantly within posterior involvement including parieto-occipital lobes and splenium of the corpus callosum; about 15% of patients have an anterior pattern of atypical involvement.^{1,2} Frontal involvement can be seen in X-ALD, and this imaging feature can expedite the diagnosis of this atypical X-ALD.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Appendix (continued)

Name	Location	Role	Contribution
Yayun Cao, MD	Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology	Author	Study concept and design, acquisition of data, analysis and interpretation of data
Bitao Bu, MD	Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology	Author	Study concept and design, critical revision of manuscript for intellectual content

Appendix Authors

Name	Location	Role	Contribution
Chenchen Liu, MD	Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology	Author	Study concept and design, acquisition of data, analysis and interpretation of data

References

1. Moser HW, Loes DJ, Melhem ER, et al. X-linked adrenoleukodystrophy: overview and prognosis as a function of age and brain magnetic resonance imaging abnormality: a study involving 372 patients. *Neuropediatrics* 2000;31: 227–239.
2. Loes DJ, Fatemi A, Melhem ER, et al. Analysis of MRI patterns aids prediction of progression in X-linked adrenoleukodystrophy. *Neurology* 2003;61: 369–374.

Neurology®

Teaching NeuroImages: Frontal lobe involvement in adult-onset cerebral X-linked adrenoleukodystrophy

Chenchen Liu, Yayun Cao and Bitao Bu

Neurology 2019;93:e1326-e1327

DOI 10.1212/WNL.00000000000008181

This information is current as of September 24, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/93/13/e1326.full
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/93/13/e1326.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://n.neurology.org/cgi/collection/all_genetics
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.