

Ventricular volume expansion in presymptomatic genetic frontotemporal dementia

Tamara P. Tavares, HBSc, Derek G.V. Mitchell, PhD, Kristy Coleman, MSc, et al., on behalf of the Genetic FTD Initiative, GENFI

Cite as: *Neurology*® 2019;93:e1699-e1706. doi:10.1212/WNL.0000000000008386

Correspondence

Dr. Finger
elizabeth.finger@lhsc.on.ca

Study objective and summary result

This study aimed to characterize the timecourse of ventricular volume expansion in persons carrying genetic mutations associated with frontotemporal dementia (FTD), and found that ventricular volume expansion is greater in presymptomatic mutation-carriers than in noncarriers who are biologically related to symptomatic mutation-carriers.

What is known and what this paper adds

Ventricular expansion is a consistently observed feature of genetic FTD at the symptomatic stage. This investigation shows that it can also be observed during presymptomatic stages.

Participants and setting

This investigation included 46 presymptomatic carriers of FTD-associated mutations in *MAPT*, *PGRN*, or *C9orf72* and 56 noncarriers. These participants were all first-degree relatives of symptomatic mutation carriers. These individuals participated in the Genetic FTD Initiative (GENFI), which recruited participants through 12 sites in Canada, Sweden, Italy, the UK, and the Netherlands.


Design, size, and duration

The participants underwent baseline and 1-year follow-up assessments that included volumetric T1-weighted MRI scans. Total ventricular volumes were calculated by summing the volumes of the left and right lateral ventricles, the third ventricle, and the fourth ventricle, and they were expressed as percentages of total intracranial volumes. Linear mixed models were used for between-group comparisons of ventricular expansion timecourses. For each participant, an expected age at symptom onset was calculated as the difference between the participant's age and the mean age at symptom onset within the participant's family.

Main results and the role of chance

Relative to the noncarriers, the presymptomatic carriers had greater annualized rates of ventricular expansion (model estimate \pm standard error, 0.35 ± 0.15 ; $p = 0.02$). Between-group differences in ventricular volumes were observed 4 years before the expected ages at symptom onset ($p = 0.04$).

Figure Total ventricular volumes plotted against years from expected symptom onset in the presymptomatic carriers (blue) and the noncarriers (red)



The dashed black line indicates the point at which between-group differences arose.

Bias, confounding, and other reasons for caution

This investigation's sample size was limited for some subgroup analyses. This investigation relied on predicted ages at onset instead of observed ages at onset.

Generalizability to other populations

This investigation's reliance on data from high-income Western countries may limit the generalizability of the results.

Study funding/potential competing interests

This work was funded by the UK Medical Research Council, the Italian Ministry of Health and the Canadian Institutes of Health Research as part of a Centres of Excellence in Neurodegeneration grant, a Canadian Institutes of Health Research operating grant (MOP 327387), and funding from the Weston Brain Institute to authors M.M and E.F. The authors report no competing interests. Go to Neurology.org/N for full disclosures.

A draft of the short-form article was written by M. Dalefield, a writer with Editage, a division of Cactus Communications. The authors of the full-length article and the journal editors edited and approved the final version.

Neurology®

Ventricular volume expansion in presymptomatic genetic frontotemporal dementia

Tamara P. Tavares, Derek G.V. Mitchell, Kristy Coleman, et al.

Neurology 2019;93:e1699-e1706 Published Online before print October 2, 2019

DOI 10.1212/WNL.00000000000008386

This information is current as of October 2, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/93/18/e1699.full
References	This article cites 22 articles, 1 of which you can access for free at: http://n.neurology.org/content/93/18/e1699.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/93/18/e1699.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Frontotemporal dementia http://n.neurology.org/cgi/collection/frontotemporal_dementia Volumetric MRI http://n.neurology.org/cgi/collection/volumetric_mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

