Teaching NeuroImages: Pericallosal curvilinear lipoma

Yao Fu, MD, PhD, Xiangyang Fang, MD, PhD,* Yanxin Li, MS,* and Xingli Zhao, MS

Neurology® 2019;93:e212-e213. doi:10.1212/WNL.00000000000007762

Correspondence

Dr. Zhao
zhao_xing_li@163.com

Figure Cranial MRI

Sagittal T1-weighted MRI demonstrates an interhemispheric slender hyperintense mass over the corpus callosum (A), with signal attenuation on fat suppression sequence (B).

A 27-year-old woman presented to the neurosurgery department with a history of an incidental finding of an intracranial lesion. Neurologic examination was unremarkable. Cranial MRI revealed an interhemispheric lipoma over the corpus callosum (figure). A diagnosis of pericallosal curvilinear lipoma was made. Intracranial lipoma is a rare congenital malformation.¹ Pericallosal region is the most common location of intracranial lipomas. Pericallosal lipoma can be divided into 2 morphologic subtypes: tubulonodular and curvilinear.² The latter is mostly asymptomatic, occasionally presenting with seizures and headache, and has a low incidence of other accompanying anomalies. In this patient, surgical resection was not necessary, and regular follow-up was recommended.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

References

1. Rajan DS, Popescu A. Corpus callosum lipoma. *Neurology* 2012;78:1366.
2. Yildiz H, Hakyemez B, Koroglu M, et al. Intracranial lipomas: importance of localization. *Neuroradiology* 2006; 48:1–7.

MORE ONLINE

→Teaching slides

links.lww.com/WNL/A917

*These authors contributed equally to this work.

From the Department of Neurosurgery, China-Japan Union Hospital of Jilin University, Changchun, Jilin Province, China.

Go to Neurology.org/N for full disclosures.

Appendix Authors

Name	Location	Role	Contribution
Yao Fu, MD, PhD	Jilin University, Changchun, China	Author	Designed and conceptualized study, analyzed the data, drafted the manuscript for intellectual content
Xiangyang Fang, MD, PhD	Jilin University, Changchun, China	Author	Drafting and revision of the manuscript
Yanxin Li, MS	Jilin University, Changchun, China	Author	Drafting and revision of the manuscript
Xingli Zhao, MS	Jilin University, Changchun, China	Author	Designed and conceptualized study, analyzed the data, drafted the manuscript for intellectual content

Neurology®

Teaching NeuroImages: Pericallosal curvilinear lipoma

Yao Fu, Xiangyang Fang, Yanxin Li, et al.

Neurology 2019;93:e212-e213

DOI 10.1212/WNL.0000000000007762

This information is current as of July 8, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/93/2/e212.full
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/93/2/e212.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.