



Abstracts

Articles appearing in the August 2019 issue

Congenital myopathies in the adult neuromuscular clinic: Diagnostic challenges and pitfalls

Objective To investigate the spectrum of undiagnosed congenital myopathies (CMs) in adults presenting to our neuromuscular clinic and to identify the pitfalls responsible for diagnostic delays.

Methods We conducted a retrospective review of patients diagnosed with CM in adulthood in our neuromuscular clinic between 2008 and 2018. Patients with an established diagnosis of CM before age 18 years were excluded.

Results We identified 26 patients with adult-onset CM and 18 patients with pediatric-onset CM who were only diagnosed in adulthood. Among patients with adult onset, the median age at onset was 47 years, and the causative genes were *RYR1* (11 families), *MYH7* (3 families), and *ACTA1* (2 families), and *SELENON*, *MYH2*, *DNM2*, and *CACNA1S* (1 family each). Of 33 patients who underwent muscle biopsy, only 18 demonstrated histologic abnormalities characteristic of CM. Before their diagnosis of CM, 23 patients had received other diagnoses, most commonly non-neurologic disorders. The main causes of diagnostic delays were mildness of the symptoms delaying neurologic evaluation and attribution of the symptoms to coexisting comorbidities, particularly among pediatric-onset patients.

Conclusions CMs in adulthood represent a diagnostic challenge, as they may lack the clinical and myopathologic features classically associated with CM. Our findings underscore the need for a revision of the terminology and current classification of these disorders.

[NPub.org/NG/9307a](https://pubmed.ncbi.nlm.nih.gov/3193070/)

Human *GABRG2* generalized epilepsy: Increased somatosensory and striatthalamic connectivity

Objective To map fMRI connectivity within and between the somatosensory cortex, putamen, and ventral thalamus in individuals from a family with a GABAergic deficit segregating with febrile seizures and genetic generalized epilepsy.

Methods We studied 5 adults from a family with early-onset absence epilepsy or febrile seizures and a GABA_A receptor subunit $\gamma 2$ pathogenic variant (*GABRG2*[R43Q]) vs 5 age-matched controls. We infer differences between participants with the *GABRG2* pathogenic variant and controls in resting-state fMRI connectivity within and between the somatosensory cortex, putamen, and ventral thalamus.

Results We observed increased fMRI connectivity within the somatosensory cortex and between the putamen and ventral thalamus in all individuals with the *GABRG2* pathogenic variant compared with controls. Post hoc analysis showed less pronounced changes in fMRI connectivity within and between the primary visual cortex and precuneus.

Conclusions Although our sample size was small, this preliminary study suggests that individuals with a *GABRG2* pathogenic variant, raising risk of febrile seizures and generalized epilepsy, display underlying increased functional connectivity both within the somatosensory cortex and in striatthalamic networks. This human network model aligns with rodent research and should be further validated in larger cohorts, including other individuals with generalized epilepsy with and without known GABA pathogenic variants.

[NPub.org/NG/9307b](https://pubmed.ncbi.nlm.nih.gov/3193071/)



Most-Read Articles

As of June 12, 2019

KCNQ2 encephalopathy: Features, mutational hot spots, and ezogabine treatment of 11 patients

J.J. Millichap, K.L. Park, T. Tsuchida, et al. 2016;2:e96. doi.org/10.1212/NXG.0000000000000096

CHCHD10 variant p.(Gly66Val) causes axonal Charcot-Marie-Tooth disease

M. Auranen, E. Ylikallio, M. Shcherbii, et al. 2015;1:e1. doi.org/10.1212/NXG.0000000000000003

The Clinical Outcome Study for dysferlinopathy: An international multicenter study

E. Harris, C.L. Bladen, A. Mayhew, et al. 2016;2:e89. doi.org/10.1212/NXG.0000000000000089

Mendelian randomization shows a causal effect of low vitamin D on multiple sclerosis risk

B. Rhead, M. Bäärnhielm, M. Gianfrancesco, et al. 2016;2:e97. doi.org/10.1212/NXG.0000000000000097

A novel *DYNC1H1* mutation causing spinal muscular atrophy with lower extremity predominance

Q. Niu, X. Wang, M. Shi, Q. Jin. 2015;1:e20. doi.org/10.1212/NXG.0000000000000017

Redefining the phenotype of ALS and AARS2 mutation-related leukodystrophy

R. Lakshmanan, M.E. Adams, D.S. Lynch, et al. 2017;3:e135. doi.org/10.1212/NXG.0000000000000135

Neurology[®]

What's happening in *Neurology*[®] *Genetics*
Neurology 2019;93;300
DOI 10.1212/WNL.0000000000007948

This information is current as of August 12, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/93/7/300.full
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

