

# Teaching Video NeuroImage: Impaired bilateral conjugate eye movements in a 48-year-old man


Mao Liu, MD, Dr med, Jing Zhang, MD, PhD, and Min Zhang, MD, PhD

*Neurology*® 2020;94:e1966-e1967. doi:10.1212/WNL.0000000000009382

**Correspondence**

Dr. M. Zhang  
mzhang@tjh.tjmu.edu.cn

**Figure** Lesion ventral to the cerebral aqueduct in the pontomesencephalic junction



Increased signal on T2 fluid-attenuated inversion recovery (C) and postcontrast enhancement (A, B, and F) is seen. Mildly increased and decreased signals on diffusion-weighted imaging (D) and apparent diffusion coefficient (E), respectively, suggest weak diffusion restriction, supporting a diagnosis of ischemic stroke in the subacute phase.

A 48-year-old man developed lightheadedness and diplopia upon awakening 10 days prior. Since then, his double vision had been continuous and stable. Upon admission, neurologic examination revealed bilateral internuclear ophthalmoplegia, abducting nystagmus, and intact convergence reflex (video). His blood pressure was 155/100 mm Hg and HbA1c was 8.7%, suggesting a new diagnosis of diabetes mellitus type 2. CT angiography showed calcified plaques of bilateral cavernous internal carotid arteries. MRI showed a lesion with hyperintensity on T2 fluid-attenuated inversion recovery, contrast enhancement, and mild diffusion restriction ventral to the cerebral aqueduct in the pontomesencephalic junction (figure), which is supplied by interpeduncular perforating branches of posterior cerebral arteries.<sup>1</sup> For this patient with hypertension and diabetes mellitus, the acute onset of symptoms and imaging findings suggest ischemic stroke due to arteriolosclerosis of perforating arteries, which should be differentiated from demyelinating diseases such as multiple sclerosis. Involvement of bilateral medial longitudinal fasciculus rather than oculomotor nucleus is a rare consequence of stroke.<sup>2</sup>

## MORE ONLINE

### Video

### Teaching slides

[links.lww.com/WNL/B78](https://links.lww.com/WNL/B78)

From the Department of Neurology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

Go to [Neurology.org/N](https://Neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

---

## Appendix Authors

Name	Location	Contribution
<b>Mao Liu, MD, Dr med</b>	Tongji Medical College, Huazhong University of Science and Technology, Wuhan	Designed the study; collected, analyzed, and interpreted the data; wrote and revised the manuscript
<b>Jing Zhang, MD, PhD</b>	Tongji Medical College, Huazhong University of Science and Technology, Wuhan	Analyzed and interpreted the data, revised the manuscript
<b>Min Zhang, MD, PhD</b>	Tongji Medical College, Huazhong University of Science and Technology, Wuhan	Designed the study, interpreted the data, revised the manuscript

## Study funding

No targeted funding reported.

## Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

## References

1. Marinkovic S, Milisavljevic M, Kovacevic M. Interpeduncular perforating branches of the posterior cerebral artery: microsurgical anatomy of their extracerebral and intracerebral segments. *Surg Neurol* 1986;26:349–359.
2. Frohman TC, Galetta S, Fox R, et al. Pearls & Oy-sters: the medial longitudinal fasciculus in ocular motor physiology. *Neurology* 2008;70:e57–e67.

# Neurology®

## Teaching Video NeuroImage: Impaired bilateral conjugate eye movements in a 48-year-old man

Mao Liu, Jing Zhang and Min Zhang

*Neurology* 2020;94:e1966-e1967 Published Online before print April 21, 2020

DOI 10.1212/WNL.0000000000009382

**This information is current as of April 21, 2020**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://n.neurology.org/content/94/18/e1966.full">http://n.neurology.org/content/94/18/e1966.full</a>
<b>References</b>	This article cites 2 articles, 1 of which you can access for free at: <a href="http://n.neurology.org/content/94/18/e1966.full#ref-list-1">http://n.neurology.org/content/94/18/e1966.full#ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>All Cerebrovascular disease/Stroke</b> <a href="http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke">http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke</a> <b>Diplopia (double vision)</b> <a href="http://n.neurology.org/cgi/collection/diplopia_double_vision">http://n.neurology.org/cgi/collection/diplopia_double_vision</a> <b>MRI</b> <a href="http://n.neurology.org/cgi/collection/mri">http://n.neurology.org/cgi/collection/mri</a> <b>Nystagmus</b> <a href="http://n.neurology.org/cgi/collection/nystagmus">http://n.neurology.org/cgi/collection/nystagmus</a> <b>Ocular motility</b> <a href="http://n.neurology.org/cgi/collection/ocular_motility">http://n.neurology.org/cgi/collection/ocular_motility</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.neurology.org/about/about_the_journal#permissions">http://www.neurology.org/about/about_the_journal#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://n.neurology.org/subscribers/advertise">http://n.neurology.org/subscribers/advertise</a>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

