Cavitating leukodystrophy as a manifestation of cerebral involvement in MFN2 neuropathy

Ranjith Kumar Manokaran, MD, DM, Harsha Vardhan Mahalingam, MD, DNB, PDF, and Deepti Kewalramani, MD

Neurology® 2020;95:42-43. doi:10.1212/WNL.0000000000009746

Correspondence

Dr. Mahalingam
harsha.mahalingam@gmail.com

A 16-year-old girl presented with progressive bilateral lower limb weakness for 3 years. Examination revealed tongue fasciculations, muscle wasting, and extensor plantar responses. Nerve conduction studies revealed motor-sensory axonal polyneuropathy. MRI brain showed multifocal cavitating white matter disease with diffusion restriction (figures 1 and 2). Exome sequencing revealed heterozygous missense variation c.775C>T (p.Arg259Cys) in exon 8 of the *MFN2* gene, pathogenic for Charcot-Marie-Tooth disease 2A (CMT2A).

Dominant mutations of *MFN2* (encoding mitochondrial protein mitofusin-2) cause a disorder of mitochondrial DNA maintenance¹ resulting in axonal sensorimotor neuropathy.

Figure 1 MRI brain: T2-weighted and T2-fluid-attenuated inversion recovery (FLAIR)-weighted images

Axial T2-weighted images (A–C) show discrete hyperintense foci in periventricular and deep cerebral white matter. Axial T2-FLAIR images (D and E) show suppression of signal within the lesions suggesting cystic nature (white arrows). A nonsuppressed lesion (F) was seen in the left temporal lobe (black arrow).

From the Division of Paediatric Neurology (R.K.M.) and Department of Radiology (H.V.M.), Sri Ramachandra Institute of Higher Education and Research (SRIHER), Chennai; and Department of Pediatric Neurology (D.K.), BJ Wadia Children Hospital, Mumbai, India.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Figure 2 MRI brain: diffusion-weighted imaging

Axial diffusion-weighted images (A–C) and corresponding apparent diffusion coefficient maps (D–F) show diffusion restriction along the walls of some cystic lesions (white arrows) and solid diffusion restriction in the left temporal lobe lesion (black arrows). Spinal cord was normal (images not shown).

Nonspecific white matter alterations are reported in few patients with CMT2A.² Multifocal cavitating leukodystrophy may be seen rarely.

Study funding

No targeted funding reported.

Disclosure

The authors report no relevant disclosures. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

References

1. Larrea D, Pera M, Gonnelli A, et al. MFN2 mutations in Charcot-Marie-Tooth disease alter mitochondria-associated ER membrane function but do not impair bioenergetics. *Hum Mol Genet* 2019;28:1782–1800.
2. Brockmann K, Dreha-Kulaczewski S, Dechent P, et al. Cerebral involvement in axonal Charcot-Marie-Tooth neuropathy caused by mitofusin2 mutations. *J Neurol* 2008; 255:1049–1058.

Appendix Authors

Name	Location	Contribution
Ranjith Kumar Manokaran, MD, DM	Division of Paediatric Neurology, Sri Ramachandra Institute of Higher Education and Research (SRIHER), Chennai, India	Conceptualization, data collection, manuscript editing
Harsha Vardhan Mahalingam, MD, DNB, PDF	Department of Radiology, Sri Ramachandra Institute of Higher Education and Research (SRIHER), Chennai, India	Data collection, manuscript drafting
Deepti Kewalramani, MD	Department of Pediatric Neurology, BJ Wadia Children Hospital, Mumbai, India	Data collection, manuscript drafting

Neurology®

Cavitating leukodystrophy as a manifestation of cerebral involvement in MFN2 neuropathy

Ranjith Kumar Manokaran, Harsha Vardhan Mahalingam and Deepti Kewalramani

Neurology 2020;95;42-43 Published Online before print June 10, 2020

DOI 10.1212/WNL.00000000000009746

This information is current as of June 10, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/95/1/42.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/95/1/42.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://n.neurology.org/cgi/collection/all_genetics Leukodystrophies http://n.neurology.org/cgi/collection/leukodystrophies Mitochondrial disorders http://n.neurology.org/cgi/collection/mitochondrial_disorders Mitochondrial disorders; see Genetics/Mitochondrial disorders http://n.neurology.org/cgi/collection/mitochondrial_disorders_see_genetics-mitochondrial_disorders MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.